



PRINCIPES

Journal of The International Palm Society

October 1984
Vol. 28, No. 4

THE INTERNATIONAL PALM SOCIETY

A nonprofit corporation engaged in the study of palms and the dissemination of information about them. The Palm Society is international in scope with world-wide membership, and the formation of regional or local chapters affiliated with The Palm Society is encouraged. Please address all inquiries regarding membership or information about the society to The Palm Society, Inc., P.O. Box 368, Lawrence, Kansas 66044, U.S.A.

PRESIDENT: Mr. Allan Bredeson, 2347 Peppermint Lane, Lemon Grove, California 92045.

VICE PRESIDENT: Mr. Edward M. McGhee, 1325 East Lake Drive, Fort Lauderdale, Florida 33136.

SECRETARY: Mr. James Mintkin, P.O. Box 27, Forestville, California 95436.

TREASURER: Mr. Ross Wagner, 4943 Queen Victoria Road, Woodland Hills, California 91364.

DIRECTORS: 1982-1986: Mr. Donn Carlsmith, Hawaii; Mr. Richard Douglas, California; Dr. Fred Essig, Florida; Mr. Don Evans, Florida; Dr. Dennis Johnson, Texas; Mr. Frank Ketchum, California; Mr. Jim Mintkin, California; Mr. Robert E. Paisley, Australia; Dr. William Theobald, Hawaii. 1984-1988: Mrs. Teddie Buhler, Florida; Dr. T. Anthony Davis, India; Mr. Garrin Fullington, N. California; Mr. Bill Gunther, S. California; Mr. Rolf Kyburz, Qld., Australia; Mrs. Lynne McKamey, Texas; Dr. Robert Read, Maryland. Others: Dr. John Dransfield, London; Mrs. Lois Rossten, California; Mrs. Pauleen Sullivan, California; Dr. Natalie Uhl, New York.

ADVISORY COUNCIL: Mr. Nat J. De Leon, Florida; Mr. Kenneth C. Foster, California; Dr. Walter H. Hodge, Florida; Dr. Jerome P. Keuper, Florida; Mr. Myron Kinnach, California; Mr. Eugene D. Kitzke, Wisconsin; Dr. John Popenoe, Florida; Mr. Dent Smith, Florida; Dr. U. A. Young, Florida; Mrs. Lucita H. Wait, Florida.

BOOKSTORE: Mrs. Pauleen Sullivan, 3616 Mound Avenue, Ventura, California 93003.

SEED BANK: Mrs. Lois Rossten, Seed Distributor, 6561 Melbourne Dr., Huntington Beach, California 92647.

PRINCIPES

EDITORS: Dr. Natalie W. Uhl, 467 Mann Library, Ithaca, N.Y. 14853. Dr. John Dransfield, The Herbarium, Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3AB England.

ASSOCIATE EDITOR: Dr. Dennis Johnson, 311 Stanford St., Hyattsville, Maryland 20783.

FIELD EDITORS: Mr. DeArmand Hull, Mr. James Mintkin, Mr. Ralph Velez.

Manuscripts for PRINCIPES, including legends for figures and photographs, must be typed double-spaced on one side of 8½ × 11 bond paper and addressed to Dr. Natalie W. Uhl for receipt not later than 90 days before date of publication. Authors of one page or more of print are entitled to six copies of the issue in which their article appears. Additional copies of reprints can be furnished only at cost and by advance arrangement.

Contents for October

A Magnificent New Palm from Madagascar John Dransfield and Natalie W. Uhl	151
Notes on the Palms of Guinea-Bissau Dennis V. Johnson	155
<i>Halmoorea</i> , A New Genus from Madagascar, with Notes on <i>Sindroa</i> and <i>Orania</i> John Dransfield and Natalie W. Uhl	163
The Native Palms of Puerto Rico Andrew Henderson	168
Where's Joey? Melvin W. Sneed	173
Vessel Elements in Roots of Young Palms Monoranjan Ghose	179
Features	
Palm Literature	162, 186
Classified	167
Letters	172
Bookstore	178
Index	188
Financial Statement	191
Palmeter	192

Cover Picture

Marojejya darianii, a new palm from mountain swamps in northeast Madagascar with Dominic Halleux. Photo by Dr. M. E. Darian. See pp. 151-154.

PRINCIPES

JOURNAL OF THE
INTERNATIONAL PALM SOCIETY
(ISSN 0032-8480)

An illustrated quarterly devoted to information about palms and published in January, April, July and October by The Palm Society, Inc.

Subscription price is \$12.00 per year to libraries and institutions. Membership dues of \$15.00 per year include a subscription to the Journal. Single copies are \$5.00 each, \$20.00 a volume. Airmail delivery \$2.50 a copy or \$10.00 a volume. The business office is located at **P.O. Box 368, Lawrence, Kansas 66044**. Changes of address, undeliverable copies, orders for subscriptions, and membership dues are to be sent to the business office.

Second class postage paid at Lawrence, Kansas

© 1984 The International Palm Society

Mailed at Lawrence, Kansas
October 26, 1984

A Magnificent New Palm from Madagascar

JOHN DRANSFIELD AND NATALIE W. UHL

*Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3AB, U.K. and
L. H. Bailey Hortorium, Cornell University, Ithaca, NY 14853, U.S.A.*

Dr. Mardy E. Darian, tireless and enthusiastic palm collector, has visited Madagascar several times in recent years seeking rare and unusual palms to introduce into his unparalleled collection at Vista, California. Collecting palms in Madagascar is fraught with difficulties, yet the palm flora is exceptionally exciting. After several false leads, in 1982 Mardy struggled to mountain swamps northwest of Maroansetra with the encouragement of Palm Society member Alfred Razafindratsira and the help of Dominic Halleux and found a truly magnificent, entire-leaved palm (see Cover). He returned with tales of this spectacular palm and tantalizing photographs. A second trip provided seeds and leaf fragments and a third trip pieces of old inflorescences.

Initially it appeared that the palm represented a new genus but as fragments accumulated, the affinities of the palm with *Marojejya insignis*, described by Humbert (1955) from the nearby Marojejy Massif, became more and more evident. Finally, in July 1984 we were able to compare the fragments collected by Mardy with the superb complete material of *Marojejya insignis* collected by Dr. H. E. Moore, Jr. in 1971. There seems no doubt that Mardy's palm is a new species of *Marojejya*. The palm is locally called *Ravimbe*—'Big Leaf.' We hereby honor with the specific epithet Dr. Darian's tireless efforts to discover and introduce this wonderful palm.

***Marojejya darianii* J. Dransf. & N. Uhl sp. nov.** Palma egregie insignis, foliis indivisis maximis; a *M. insigni* petiolo

carenti, vaginis auriculas maximas ferentibus, rachillis staminatis foveas profundiores ferentibus, rachillis pistillatis numerosioribus, seminibus subreniformibus, profunde-sulcatis, bene distincta. Typus: Madagascar, Maroansetra, *Darian s.n.*

Robust, solitary, monoecious, pleoanthic palm. Stem erect to 8 m tall, ca. 15–20 cm diam., when young covered in leaf bases and then appearing ca. 30 cm diam.; internodes short, bearing scattered, short, spinelike adventitious roots. Leaves massive, ca. 20–30 in the crown, held more or less stiffly erect, entire, bifid, pinnately ribbed, becoming torn by the wind; sheath with 2 conspicuous, large rounded auricles, ca. 10–12 cm wide; petiole absent; rachis ca. 15 cm wide near the base, very spongy in texture; blade ca. 7–9 m long, bifid in the apical 20–50 cm, basally the margins long decurrent to the sheath, gradually widening distally, ca. 1–1.2 m wide at the widest point, ca. $\frac{2}{3}$ the blade length from the base; margins finely serrate; adaxial surfaces of the ribs and blade glabrous, abaxial surface of ribs bearing abundant pale, floccose scales, the blade abaxially with scattered, minute, dotlike scales. Inflorescences ?unisexual. Staminate inflorescence unknown except for rachilla fragments; rachillae more or less catkinlike, length unknown, ca. 7–8 mm diam., apparently bearing flowers to the tip; rachilla bracts, 2×1.75 mm, spirally arranged, congested, horizontally inserted, joined laterally and forming pits ca. 2 mm diam., the free tips apiculate, distally pointing, the exposed part densely tomentose except at the tip, the tip



1. Dr. Darian points to the conspicuous auricle on the leaf of *Marojejya darianii*; note the absence of a petiole.




2. An old pistillate inflorescence of *Marojejya darianii*.

extending between the staminate flowers; floral bracteoles minute, only 1 seen. Staminate flowers borne in pairs, more or less symmetrical, ca. 3.5×2.0 mm; sepals 3, free, widely separated, tongue-like, ca. 2.25×1 mm, apically pointed, slightly keeled, the margins very finely serrate, 1 sepal usually slightly wider than the others; petals 3, free, valvate, more or less boat-shaped, 2.5×1.5 mm; stamens 6, filaments awl-shaped, long and slender, ca. 2.0–2.5 mm long, united at the base with the pistillode, the antesealous inserted lower than the antepetalous; anthers medifixed, ca. 2.5×1 mm, latrorse, more or less exerted at anthesis; pistillodes 3, irregularly joined, ca. 0.75 mm long. Pistillate inflorescence (Fig. 2) apparently partially concealed among the leaf bases; peduncle at least 15 cm long, more or less circular in cross-section, ca. 2 cm diam.; prophyll and peduncular bracts not available; rachis ca. 6 cm long, bearing 48 closely crowded, catkinlike rachillae;

rachillae more or less equal, ca. $10-11 \times 1.3$ cm, bearing spirally arranged bracts united laterally and basally to form pits; pits ca. 4.5 mm diam.; rachilla bracts triangular, apiculate, ca. 3×4 mm, at first apparently closing the pit, becoming reflexed; abortive staminate flowers 2, very small, concealed within the pit, lateral to the pistillate, floral bracteoles 2, broadly triangular, 2×2 mm. Pistillate flowers unknown. Mature fruit not available, said to be rounded, smooth. Seed subreniform, ca. $20 \times 12 \times 10$ mm, the surface covered with deep, broad, mostly longitudinal, anastomosing grooves; endosperm homogeneous; embryo lateral to the hilum. Germination adjacent ligular; eophyll bifid, epetiolate, with long decurrent blade margins.

MADAGASCAR. Northeast: In mountain swamps west northwest of Maroansetra, ca. 500 m altitude, *M. Darian s.n.* 1983. (Holotype BH, isotype K.)

Notes: *Marojejya darianii* is immediately distinguishable from *M. insignis* by the absence of a petiole and by the extraordinary pointed auricles on the leaf sheath (see Fig. 1). The staminate rachillae have much more pronounced pits but the flowers are almost identical to those of *M. insignis*. In the pistillate inflorescence there appear to be many more rachillae in the new palm. Perhaps the most striking difference is in the seed (Fig. 3); that of *M. darianii* is subreniform and deeply grooved, whereas that of *M. insignis* is irregularly globular and only very faintly grooved. However, the two seeds are basically similar in form and in the position of the embryo in relation to the hilum. The seedlings are easily separated; that of *M. darianii* bears epetiolate leaves, whereas that of *M. insignis* has definite petioles. These differences establish that the two palms are clearly specifically distinct, but there are many similarities such as the form of the leaf, inflorescence structure, and staminate flowers indicating that the two taxa are congeneric.



3. *Marojejya darianii*, a-c, the seed in three views; d, seed in longitudinal section; e, in transverse section. *Marojejya insignis*, f, view of seed to show the position of the embryo, lateral to the hilum, similar to that of *M. insignis* in c. Drawn by Soejatmi Dransfield.

LITERATURE CITED

HUMBERT, H. 1955. Une merveille de la nature à Madagascar. Première exploration botanique du Massif du Marojejy et de ses satellites. Mém.

Inst. Rech. Sci. Madagascar. Sér. B. Biol. Vég. 6: 1-271.

Note: *M. darianii* grows at temperatures of 50-80 F under deep shade in humus of sandy swamps where rainfall is very high. Dr. M. E. Darian, 2615 So. Santa Fe Ave., Vista, CA 92083, has seedlings of both *M. darianii* and *M. insignis* for exchange with Botanic Gardens and other collectors.

Principes, 28(4), 1984, pp. 155-162

Notes on the Palms of Guinea-Bissau

DENNIS V. JOHNSON

3311 Stanford St., Hyattsville, MD 20783, U.S.A.

Remote and lesser-known countries hold a certain fascination for the visitor because there is a sense of discovery in observing how people live and make use of their natural environment. Guinea-Bissau, wedged between Senegal and Guinea in West Africa, is such a place. Formerly Portuguese Guinea, this small nation is about equal in area to the states of Massachusetts and Connecticut.

The literature on African vegetation commonly mentions the paucity of palm genera and species in this the largest tropical area of the world (see Tomlinson 1962). Although smaller in area, tropical Asia and tropical America each has more

types of palms. The contrast is strikingly shown by the fact that Cuba has a greater diversity of palms than continental Africa (Corner 1966). But what may be lacking in diversity is, at least in West Africa, more than compensated for by the sheer numbers of palms. They represent a common element of the vegetation landscape. Most prevalent is the African oil palm (*Elaeis guineensis*), certainly as useful a plant in West Africa as the coconut is in Polynesia. This holds true for Guinea-Bissau. Of secondary importance is the African fan palm (*Borassus aethiopum*), found throughout the continent.

I had the good fortune of spending three



1. Subspontaneous oil palms amid and adjacent to rice fields near Saucunda.



2. Utensils used for simple manual extraction of palm oil on a farm near Bafatá.

months in Guinea-Bissau in 1982, working as a consultant on an agricultural development project in the central part of the country. I took the opportunity to collect some notes on palm products, particularly those from the two mentioned above.

African Oil Palm

This pinnate-leaved palm has a dual role in the Guinea-Bissau economy in that it provides a major export (kernels) and furnishes palm oil, palm wine and other products for local consumption. Oil palms are abundant in the littoral zone, on the offshore islands and in the river valleys. Because of the long association with human activities, it is impossible to determine the precise ecological niche of this palm anywhere in West Africa. For that reason it is most often described as being subsynchronous. When forests containing oil palms are cleared for agriculture, the palms are commonly left standing (Fig. 1). Over generations, and if allowed to regenerate naturally, the palms form nearly pure

stands. The palm's resistance to fire, extensively used in land clearing and for weed control, also gives it an advantage over other woody species.

Oil palms in Guinea-Bissau are of the *dura* type, which have large kernels and low to medium mesocarp content. Estimates place the area under oil palm in the country at about 100,000 ha. This figure is for natural stands; formal plantations do not exist.

Apart from a single industrial palm oil processing plant in Bubaque, on one of the offshore islands, the extraction of palm oil is a simple cottage industry solely for the domestic market. Palm oil is derived from the mesocarp pulp of the fruit, has a reddish-orange color because it is rich in carotene and is locally known as *che-beu*. The kernel contains a clear oil designated in the trade as palm kernel oil. Figure 2 shows some of the utensils used in the manual extraction process. Fruit bunches are harvested and transported to a local processing site. The fruits are removed from the bunches, loaded into



3. Oil palm kernels drying in the sun wreathed by empty fruit bunches on a farm near Bafatá.


large baskets such as the one in the foreground of the photograph, and dumped into used oil drums of boiling water. After being cooked for a few hours, the fruits are removed from the water, pounded to break up the pulp, and then returned to the boiling water. The oil separates from the pulp and floats to the top where it is skimmed off. The pans in the photograph are used to transport the pulp and the palm oil. The crude oil is filtered of impurities and sold in various-sized recycled glass bottles. This is the traditional cooking oil of the country and is also said to have cosmetic and medicinal applications.

Most of the pulp having been removed during the extraction of palm oil, the remaining seeds are spread in the sun to dry and then more commonly cracked by hand to remove the kernels. Mechanized shelling by means of a small portable gasoline-powered machine is now being done to an undetermined degree in Guinea-Bissau. The extracted kernels are dried in the sun (Fig. 3) and exported. There is not the industrial capacity within the


country to express the palm kernel oil. Production of palm kernels in 1982 was estimated to be 10,000 mt; that of palm oil 5,500 mt (FAO 1983).

The tapping of oil palms for palm wine is commonplace in Guinea-Bissau and throughout West Africa. Palm wine is the usual designation for the sap which ferments very quickly into a weak alcoholic beverage. This is equivalent to the palm toddy of Asia.

Oil palms selected for tapping must be cleared of dead leaves to permit easy access to the male inflorescences. A small incision is made in an unopened inflorescence and a receptacle positioned below it to collect the sap. A small funnel is made from a palm leaflet and placed in the neck of the bottle to direct the dripping sap. The man pictured in Figure 4 is ready to climb a palm with the aid of a beltlike device and empty the glass bottles of palm wine. The more traditional receptacle in Guinea-Bissau is a clay bottle (*moringo*) (Fig. 5). The bottle rests in a sling made of oil palm leaf fiber. The two loose ends



4. Tapper preparing to climb an oil palm near Saucunda.



5. Clay bottle suspended in its sling made of oil palm leaf fiber for palm wine collection near Bafatá.



6. Clay bottles attached to an oil palm for wine collection near Bafatá.



7. Oil palm trunks used for bridge construction near Jabicunda.

of the sling are fastened around the inflorescence to hold the bottle in place.

The beltlike device for climbing oil palms in Guinea-Bissau (Fig. 4) deserves further elaboration. Just as a paratrooper folds his own parachute, tradition has it that the palm wine tapper fashions his own belt, which is made from a fresh oil palm leaf. The midrib is stripped and bent to give it an elliptical shape. One end of the midrib is cut thin and flexible with a thick piece left at the extreme end to facilitate tying it to the opposite end where the midrib fibers have been separated and braided into a rope about 60 cm long. These ends are on the left side of the belt. Where the belt comes in contact with the trunk it often is wrapped with extra fiber to reduce wear. To climb a tree the tapper loops one end of the belt around the trunk and the other around his lower back, and secures the two ends. Grasping the sides of the belt, he leans forward and jumps the belt

up the trunk a short distance. Then, leaning back against the belt, walks an equal distance up the rough trunk. The knot is secure enough that when he reaches the crown the tapper can have both hands free for work. Anyone who has seen an electrician ascend a wooden utility pole using a leather belt and spikes attached to his shoes can visualize the process of collecting palm wine. A belt made from an oil palm leaf midrib lasts about one month.

On one of my field trips I was able to borrow a belt and, to the great amusement of those assembled, made a shaky but successful ascent of an oil palm. Quite taken by the ingenuity of the belts, I later had one made. It, along with a clay bottle in a sling, occupies a prominent place on the wall of my office.

Palm wine yields vary considerably, but I was told that 10–15 liters per day per tree was about average. The oil palms are



8. African fan palms in the village of Nhacra.

tapped only during the dry season when the sap yield is sufficient to justify the labor. Inflorescence tapping decreases the tree's fruit yield, but apparently does not do any permanent damage. Palm wine has a milky appearance because of its high yeast content. The taste is pleasant. I sampled it on several occasions under different circumstances and found it best just after being collected (fermentation occurs within the receptacle on the tree) and at ambient temperature. Within a few hours palm wine becomes unpleasantly bitter. This beverage is popular among non-Muslims in the rural areas and is a good source of vitamin B complex. In 1982 a liter of palm wine cost the equivalent of 40 cents. It is not marketed in any organized fashion. Because of its high yeast content, palm wine is used in making leavened bread.

In Guinea-Bissau there is no current

program to produce commercially and market palm wine, although the technology has been developed in Nigeria. There, the bottling and preservation of palm wine was investigated by Levi and Oruche (1957); a study of palm wine production was carried out by Tuley (1965); the Nigerian Institute for Oil Palm Research conducted experiments on tapping palms in the late 1960s and early 1970s; and the potential for improving traditional oil palm wine production was the subject of a recent article by Okereke (1982).

Other uses of the oil palm in Guinea-Bissau include weaving coarse baskets and mats from the leaves. These also are employed for thatching and to make fences. Small bridges are constructed with oil palm trunks (Fig. 7). The palm heart is eaten when trees are felled. Weaverbirds like the oil palm and use the leaflets

to build their distinctive suspended nests. The oil palm is normally unbranched, but if some physical damage causes it to fork, such a tree in West Africa is thought to have taboo properties (Gledhill 1972).

African Fan Palm

In Guinea-Bissau, this attractive palm has the vernacular name *cibe*. Although it is not a source of oil, it furnishes other products similar to those from the oil palm. The classic savanna palm of Africa, it can be found growing in close proximity to the oil palm in Guinea-Bissau. The number of African fan palms within the country has been reduced significantly through a combination of land clearing, cutting the tree for its wood and insect pests which have become more severe as a result of upsetting the ecological balance by removal of the natural forest (Castel-Branco and Tordo 1956). A few relatively young palms are shown in Figure 8.

For purposes of weaving and thatching, African fan palm leaves are the preferred type. Quality hats, baskets and purses are woven from the leaves and command the highest prices on local markets. The entire leaf is employed in thatching, frequently as the supporting layer for the traditional grass roofs. I have seen individuals carrying freshly-cut leaves as rain umbrellas and the leaves spread over charcoal to keep it from getting wet.

The African fan palm bears green fruits which are about the size of an orange. When immature, the soft endosperm can be eaten raw and the mesocarp pulp sliced and cooked. The pulp of the ripe fruit is eaten fresh. When felled, the palm heart is extracted from the trunk and eaten. These palms also are tapped for palm wine in the same manner as described for the oil palm. Exploitation of this palm is not as widespread as it once was, due to their reduced numbers. I saw relatively few tall

mature trees, possibly because it is they which furnish the best wood.

Other Palms

At least three other palm genera may be native to Guinea-Bissau. Raffia palms (*tara*) grow along the coast according to Mota (1954). Based upon the distribution of *Raphia* species provided by Russell (1964), *R. palma-pinus* and *R. sudanica* may be represented. I was unable to investigate utilization of raffia palms; however, in Nigeria, the palms are tapped for wine and, in the 19th century, oil was extracted from the fruit pulp on a small scale. Raffia oil is similar in quality to palm oil and has the same uses (Otedoh 1974).

The coastal forests of Guinea-Bissau also probably harbor climbing rattans, although I have no confirmation of this. The geographic range of *Laccosperma* (*Ancistrophyllum*) *secundiflorum* and *Calamus deeratus* reported by Irvine (1961) suggests that they may be found within the country.

Coconuts have been in West Africa since their 16th-century introduction by the Portuguese, and probably reached Guinea-Bissau as early as any place in the region. This palm is cultivated informally in the coastal region of the country. Husked coconuts are sold in local markets and copra made for export. Production of coconuts in 1982 was estimated to be 25,000 mt and copra production 5,000 mt (FAO 1983).

Acknowledgments

I wish to thank Mário Samba Embaló for helping arrange visits to areas of palm oil processing and palm wine collection, and Kit Macy for obtaining elusive documents about the oil palm.

LITERATURE CITED

- CASTEL-BRANCO, A. J. F. AND G. C. TORDO. 1956.
Acerca do equilíbrio bioecológico dos povoa-

- mentos de "cibes" *Borassus* ssp. na Guiné Portuguesa. Estudos, Ensaios e Documentos 26, 68 pp. Junta de Investigações do Ultramar, Lisbon.
- CORNER, E. J. H. 1966. The natural history of palms. University of California Press, Berkeley.
- FAO. 1983. FAO production yearbook 1982. vol. 36. Food and Agriculture Organization of the United Nations, Rome.
- GLEDHILL, D. 1972. West African trees. Longman, London.
- IRVINE, F. R. 1961. Woody plants of Ghana. Oxford University Press, London.
- LEVI, S. S. AND C. B. ORUCHE. 1957. The preservation and bottling of palm wine. Research Report No. 1, Federal Institute of Industrial Research, Lagos.
- MOTA, A. T. DA. 1954. Guiné Portuguesa. 2 vols. Agência Geral do Ultramar, Lisbon.
- OKEREKE, O. 1982. The traditional system of oil palm wine production in Igbo Eze local government area of Anambra state of Nigeria. *Agricultural Systems* 9: 239-253.
- OTEDOH, M. O. 1974. Raphia oil: its extraction, properties and utilization. *Journal Nigerian Institute Oil Palm Research* 5(19): 45-49.
- RUSSELL, T. A. 1964. The *Raphia* palms of West Africa. *Kew Bulletin* 19: 173-196.
- TOMLINSON, P. B. 1962. Palms of Africa. *Principes* 6: 96-103.
- TULEY, P. 1965. Studies on the production of wine from the oil palm. *Journal Nigerian Institute Oil Palm Research* 4(15): 284-289.

Principes, 28(4), 1984, p. 162

PALM LITERATURE

MCCOY, R. E. (ed.) Lethal Yellowing of Palms. Bulletin 834, 100 pp. Institute of Food and Agricultural Sciences, University of Florida, Gainesville. 1983.

Several articles on lethal yellowing have appeared in *Principes* since an epidemic of the disease occurred in southern Florida in the early 1970s. A serious threat to plantations of tall varieties of coconut throughout the world, lethal yellowing also attacks more than two dozen other economic and ornamental palms.

This book draws together the results of a decade of research into the origin, cause and control of lethal yellowing. Ten scientists who worked on the problem in Florida are listed as coauthors. The book is divided into seven chapters and handsomely illustrated with 34 figures, including 21 color photographs. It contains an extensive bibliography of 176 references.

Strong evidence now supports the thesis that a planthopper (*Myndus crudus*) is the insect vector of the disease. Recommendations for control include eradication of diseased palms, antibiotic treatment of infected trees while resistant palms are established to replace them, and containment of the disease. The latter is not encouraging as lethal yellowing was identified in the Yucatan Peninsula of Mexico in 1982. However, any success in slowing the spread will permit more research to be completed and allow for substitution of resistant palms in threatened areas.

The authors are to be congratulated on this excellent study of such a serious palm disease.

DENNIS JOHNSON

Halmoorea, A New Genus from Madagascar, with Notes on Sindroa and Orania

JOHN DRANSFIELD AND NATALIE W. UHL

*Herbarium, Royal Botanic Gardens, Kew, Richmond, Surrey TW9 3AB, U.K. and
L.H. Bailey Hortorium, Cornell University, Ithaca, NY 14853, U.S.A.*

One of the major difficulties in the elucidation of the limits of palm genera from Madagascar lies in the great scarcity of good material from the island. The type specimens are frequently inadequate and the protologues do not even match the specimens they purport to describe. The monotypic genus *Sindroa* exemplifies these problems. Jumelle (1933) based *Sindroa* on a collection made by Perrier (11937) in the mountains of the Masoala Peninsula at Marambo. *S. longisquama* was described as being dioecious and bearing acute leaflets, yet the inflorescences of the type specimen show clear signs of bearing flowers of both sexes and the leaflets are clearly praemorse (attached to the type is a note in Perrier's hand to the effect that the leaflets are not acute because they had been eaten in the bud by rhinoceros beetle!). With this inauspicious beginning *Sindroa* remained almost unknown.

In April 1971, H. E. Moore searched for *Sindroa* on the Masoala Peninsula and collected (9921) a handsome distichous leaved palm, bearing some of the characters of *Sindroa* and furthermore called "sindroa" by the local people. Thus it seemed that *Sindroa* had been recollected. However some features of the distichous palm did not match the type of *S. longisquama*; in particular the leaflets of the distichous palm bore pale grey brown rather than dark reddish brown ramenta. Then in December 1972, Moore revisited the Masoala Peninsula and discovered and

collected (10115) a palm with spirally arranged leaves which also seemed to be congeneric with *Sindroa*. The leaflets of the spirally arranged palm bore dark reddish brown ramenta as in the type of *S. longisquama*, and, although the type bears flowers and Moore 10115 fruit, the two collections are clearly conspecific. Moore (in his field notebook) noted the great similarity between *Sindroa* and Asiatic *Orania* even in the rather unusual seedling; on the basis of these observations *Sindroa* was placed next to *Orania* in the "Major Groups of Palms and Their Distribution" (Moore 1973). In his field notes, it appears that Hal Moore was not certain whether the distichous palm was specifically distinct from *Sindroa longisquama*.

Bernardi recollected the distichous palm (*Bernardi* 14450) and described and illustrated it in a paper as *Sindroa longisquama* (Bernardi 1974). Unfortunately the copious fruiting material collected by Bernardi has not yet been distributed from Tananarive.

In reevaluating *Sindroa* for Genera Palmarum we have come to the conclusion that the only difference separating *S. longisquama* from the rest of the genus *Orania* (excluding *O. appendiculata*) is the presence of ramenta on the undersurface of the leaflet midribs. This vegetative character is too trivial for the separation of genera and thus we believe *Sindroa* to be congeneric with *Orania*. The rather wide geographical disjunction is paralleled by many angiosperms. *Sindroa* is thus





1. Crown of *Halmoorea trispatha*, Moore 9921.

formally reduced to synonymy in *Orania* and the combination *O. longisquama* published.

***Orania longisquama* (Jumelle) J. Dransf. & N. Uhl, comb. nov.**

Sindroa longisquama Jumelle in Ann. Mus. Col. Marseille sér. 5.1 (1): 11. 1933; Jumelle & Perrier de la Bâthie, Fl. Madagascar 30: 160. 1945. Type: Madagascar, Masoala, Perrier 11937 (P, holo.!).

The distichous palm collected by Hal Moore as *Sindroa*, however, is clearly different from *Orania longisquama*. The inflorescence bears a prophyll and two large peduncular bracts rather than a prophyll and a single large peduncular bract as in *Orania*. The staminate flower has three large blunt or subacute, free, imbricate sepals (rarely two of them briefly connate), and bears 27–30 stamens, whereas in *O. longisquama* the sepals are united and acute and the stamens number 9–15. In the pistillate flower the staminodes number 12 whereas in *Orania* there are only 3–6. The distichous palm is thus abundantly distinct from *O. longisquama*. Furthermore the distichous palm does not fit within the variation found in *Orania*. We conclude that it represents a distinct genus albeit related to *Orania*. *Halmoorea* is an obvious choice of name for this extraordinary palm.




2. A. Inflorescence of *Halmoorea trispatha* in bud; p, prophyll. B. Inflorescence bud opened to show prophyll and two peduncular bracts.

***Halmoorea* J. Dransf. & N. Uhl, gen. nov.**

Palma solitaria, inermis, pleonantha, monoeca. Folium reduplicatum, pinnatum, pinnis praemorsis ramentas infra ferentibus. Inflorescentia interfoliacea trispatha, bracteis peduncularibus magnis, tubularibus, rostratis; rachillae basin triades versus apicem flores staminatus singulariter vel binatim ferentes. Flos staminatus sepalis 3, liberis, imbricatis (2 raro connatis), petalis 3, liberis, valvatis, staminibus 27–30, pistillodio carenti. Flos pistillatus sepalis 3 truncatis, basin connatis, petalis 3, liberis, valvatis, staminodiis 11–12, gynoeicio triloculato triovulato. Fructus 1, 2, vel 3-lobatus. Genus egregium Madagascariense Oraniae affine sed inflorescentia bracteis 3 (vice 2), sepalis floris staminati fere liberis et numero staminum staminodiorumque differt.

Solitary, moderate, unarmed, pleonanthic, monoecious palm. Stem erect, bare, internodes short, irregular, nodes prominent. Leaf distichous, reduplicately pin-



3. Close-up of unopened flowers from inflorescence of Figure 2.

nate, neatly abscising; sheath splitting, short, thick, woody, densely covered with brown scales, fibrous along the margins; petiole elongate, stout, adaxially channeled, abaxially rounded, with pale indument abaxially; rachis much longer than the petiole, curved, abaxially rounded, adaxially angled except at the base where grooved, with indument as the petiole; leaflets single-fold, numerous, regularly arranged in one plane, rather stiff, linear-lanceolate, obliquely praemorse, green above with prominent midnerve, 2 marginal nerves, and numerous tertiary



4. Photograph presumed to be of *Halmoorea* taken in Madagascar by Dr. M. E. Darian.

nerves, adaxially \pm glabrous, abaxially with thin gray indumentum, small punctiform scales, and irregular large gray-brown ramenta along the main veins especially near the base, transverse veinlets obscure. Inflorescence interfoliar, about as long as the petiole, branching to 3 orders; peduncle stout, covered in caducous brown scales; prophyll \pm ovate, 2 winged, persistent, margins irregular, irregularly splitting and becoming fibrous at the tip; peduncular bracts 2, tubular, inflated, beaked, both entirely enclosing the inflorescence, \pm woody, deciduous at anthesis, densely covered with irregularly mar-

gined, brown scales; rachis longer than the peduncle, brown tomentose, bearing spirally arranged, low, glabrous, coriaceous, collar-like bracts subtending first order branches; first order branches bearing few, spirally arranged, similar bracts, each subtending a 2nd order branch; 2nd order branches with a basal bare portion and few small, spirally arranged bracts subtending rachillae, all axes finely roughened; rachillae moderate, irregular, bearing up to 10 spirally arranged triads basally, paired or solitary flowers distally; rachilla bracts and bracteoles low, rounded, inconspicuous.

Staminate flowers asymmetrical, fleshy, angled; sepals 3, free, imbricate or rarely 2 united, rounded, margins somewhat irregular; petals 3, free, valvate, ovate-triangular, blunt or subacute, thickened at the tip; stamens 27–30, filaments short, slender, erect, anthers linear-oblong, bifid at the base and apex, basifixed, latrorse, connective broad; pollen \pm circular, monosulcate with finely rugulate, semitectate exine; pistillode absent. Pistillate flower only slightly larger than the staminate; sepals 3, rather large, free except basally where very briefly united, irregular, apically truncate, centrally thickened, laterally striate; petals 3, free, irregular, broadly triangular, valvate, abaxially smooth, adaxially irregularly roughened at the thickened tips; staminodes 11–12, free, threadlike, pointed; gynœcium trilobular, triovulate, irregularly obovoid, with 3 round bulges, one larger, stigmas 3, terete, ovule laterally attached, pendulous, probably hemianatropous. Fruit developing from 1, 2, or 3, carpels, rounded, bi- or trilobed, relatively large, further details not known.

Type species *H. trispatha*

Halmoorea trispatha *J. Dransf. & N. Uhl* sp. nov. Palma elata, ad 20 m; folia disticha, vaginis ferruginoso-lepidotis, petiolis ad 1 m longis, rachidibus 2 m longis, pinnae untrinque ca. 62, infra ramen-

tas pallidas ferentes. Sepala floris staminati 1.5 mm longa, petala 9×5 mm; sepala floris pistillati 5×5 mm, petala 10×8 mm; ovarium 5×8 mm, triovulatum; fructus 1, 2 vel 3-lobatus, ad ca. 6 cm latus. Typus: Madagascar, Prov. Diego Suarez, Masoala, *Moore 9921* (holotypus BH).

Moderate palm; trunk to 20 m tall, 20–35 cm dbh, swollen basally to 40 cm diam.; bark grey, weathered; internodes ca. 15 cm near the base, much shorter above, nodal scars prominent; crown with ca. 10 leaves held distichously in a fan; sheaths not forming a crownshaft, densely rusty brown scaly; petiole ca. 1 m long, ca. 4 cm diam., adaxially channeled, abaxially rounded, grey-green, covered in pale scales; rachis ca. 2 m long, abaxially rounded, adaxially angled, covered in pale indumentum; leaflets ca. 62 on each side, regularly arranged in one plane, abaxially green, adaxially grey-waxy and dotted with minute membranous brown scales, apex obliquely praemorse, midrib prominent adaxially, abaxially bearing abundant, large, lacinate ramenta, one pair of large veins prominent near the margin also bearing ramenta abaxially; transverse veinlets obscure; proximal pinnae ca. 65×1.5 cm, median pinnae ca. 75×3.5 cm, distal pinnae ca. 10×0.5 cm. Inflorescence spreading; prophyll woody, 2-keeled, green but densely rusty-scaly, to 30×10 cm; peduncular bracts 2, the first inserted 6–10 cm above the prophyll, terete, woody, inflated, beaked, brown, ca. 80×10 cm, 2nd peduncular bract inserted ca. 10 cm above the first, similar but not inflated; peduncle ca. 16–20 cm long; rachis ca. 30 cm long, bearing 10–12 branches, each subtended by an inconspicuous rachis bract; rachillae glabrous, ca. 10 cm long, probably becoming much longer at anthesis, somewhat flexuous, bearing about 10 triads basally and paired to solitary flowers distally; rachilla bracts very small, rounded; floral bracteoles very low, rounded. Staminate flowers asym-

metrical, fleshy; sepals rounded, ca. 1.5 mm long; petals broadly triangular, ca. 9×5 mm; stamen filaments very short, anthers 6×1 mm. Pistillate flowers asymmetrical; sepals thick, \pm keeled, striate laterally, ca. 5×5 mm; petals thick, ca. 10×8 mm; staminodes thread-like, ca. 2 mm; ovary 3-lobed, ca. 5×8 mm, stigmas papillose ca. 1.5 mm long. Fruit apparently 1, 2, or 3 seeded, green, the 3 seeded ca. 6 cm diam., stigmatic remains apparently basal.

Type: Madagascar, Prov. Diego Suarez, Masoala, across Onive River from Ambatobe and upriver from Ambohitralanana, 8 April 1971, *H. E. Moore 9921* (BH). Vernacular name "Sindroa."

Notes: The cabbage is said to be ined-

ible. Dr. M. E. Darian informs us that there may be two species in this genus, based on differences in trunk and seed size. Unfortunately we have no other material definitely identifiable with Hal Moore's collection. Further collections are essential.

LITERATURE CITED

- BERNARDI, L. 1974. *Notulae ad Sindroa*, genus endemicum Palmarum peninsulae Masoala, Madagascariae; cum digressionem circa Humbertiam Madagascariensem. *Candollea* 29: 163-171.
- JUMELLE, H. 1933. Nouveaux Palmiers de Madagascar. *Ann. Mus. Colon. Marseille. sér. 5.1*(1): 11.

CLASSIFIED

AVAILABLE AT THIS TIME. Seedlings of *Gronophyllum ramsayii*, *Phoenix rupicola*, *Neodypsis decaryi*, *Latania loddigesii*, *Bismarckia nobilis*, *Coccothrinax crinita*, and many others. Write for price list. RICHARD RUDY, P.O. Box 252, Winter Beach, FL 32971. Phone 305-562-1072.

DWARF RHAPIS EXCELSA, Seven green and variegated varieties available. NEW BOOK, "Secret of the Orient," a comprehensive guide to **Rhapis** palms—52 pages fully illustrated. Catalogue \$1. Book and catalogue \$5 ppd. ("Secret of the Orient" is also available for The Palm Society Bookstore). RHAPIS GARDENS—PS, P.O.D. 287, GREGORY, TX 78349.

FOR SALE. Seeds of *Jubaeopsis caffra*—\$2.50 each. PAULEEN SULLIVAN, 3616 Mound Ave., Ventura, CA 93003.

The Native Palms of Puerto Rico

ANDREW HENDERSON


The New York Botanical Garden, Bronx, New York, NY 10458, U.S.A.

Puerto Rico is one of the Antilles, an island chain curving from Florida to Venezuela, dividing the Atlantic Ocean from the Caribbean Sea. It is the easternmost of the four Greater Antilles, the others being Cuba, Jamaica and Hispaniola. It is also the smallest, about 100 miles long and 40 miles wide. The flora is distinctly Caribbean, and while the palms are not so diverse as those of Cuba, they are no less interesting. Puerto Rico has ten native species in ten genera, and half of these are endemic (Little and Wadsworth 1964; Little, Woodbury, and Wadsworth 1974).

The most widespread and abundant species is *Roystonea borinquena*, the "palma real." This palm is endemic to Puerto Rico and three small neighboring islands. It is a large and imposing palm, easily recognized at a distance by its stout grey trunk and large head of pinnate leaves. It is seen throughout the island in lowland forests, fields and planted as ornamentals. The fruits are used by farmers to feed hogs. It is said that you can tell the number of hogs a farmer owns by the number of roystoneas he has on his land, usually four palms to a hog. Although farmers are thus inclined to conserve these palms, it seems that roystoneas actually benefit from forest clearance, and can extend their range in cleared areas. There are about eleven other species of *Roystonea*, with a circum-Caribbean distribution. However, their variability, and frequency with which they are planted, means that their taxonomy is in a state of some confusion. *Roystonea borinquena* has a distinctive bulge in the trunk, and so do some of the other species. The explanation

of this is unknown, and it seems unlikely that it is entirely due to climatic changes.

The pattern of forest destruction in Puerto Rico is the same as that found in many areas throughout the Caribbean. During the last 100 years there has been extensive deforestation, leaving today only a small fraction of climax vegetation. Fortunately a well-developed conservation program has ensured that some of the most interesting vegetation types are preserved in national parks. These include subtropical dry forests, e.g., at Guanica, subtropical wet forest, e.g., at Maricao, and both



1. *Roystonea borinquena*, the most frequent palm in Puerto Rico.




2. *Thrinax morrisii*, growing in limestone rocks.

lower montane wet forest and lower montane rain forest, e.g., at Luquillo. Also secondary forests have increased in area in recent years as a result of the abandoning of marginal agricultural areas.

In the limestone hills of the island, *Thrinax morrisii* is common as an understory shrub. Typically it occurs in large populations of sterile stemless plants with a few fertile specimens with stems up to 2 m. The habitat of this species is similar to that of other related Caribbean palms, e.g., *Coccothrinax*, and consists of xeric limestone areas with a seasonal subtropical climate. The common name of *T. morrisii* is "palma de escoba," and this refers to the practice of making brooms from the leaves. This was a common industry many years ago, which then declined. Now, because of the economic recession, the practice has started up again, and brooms can be bought in local shops.



3. *Sabal causiarum*, the Puerto Rican hat plant.



4. *Goussia attenuata*, the spine-like projections on the roots, anchored in limestone rocks.




5. *Calyptronoma rivalis*, the rarest palm in Puerto Rico, growing beside a stream near San Sebastian.

Thrinax morrisii is very similar to another palmate-leaved palm, *Coccothrinax alta*, but the latter is more common in wetter limestone hills in the north of the island. The two genera can easily be separated because the bases of the petiole in *Thrinax* are split, while they are not in *Coccothrinax*.

The only other palmate-leaved palm in Puerto Rico is *Sabal causiarum*. This is another endemic species, and is found on coastal plains in the north and west of the island. Its stout grey trunks, up to 10 m, are a distinctive feature. It is known as the "palma de sombrero," although the practice of making hats from the leaves has declined over the years, with cheaper ones being made in other islands. *Sabal* is another Caribbean genus, but with wider inland extensions. Typically it inhabits lowland coastal regions, often on sandy soil. Again there is great diversity in this genus, leading to much confusion in naming.

There are two spiny pinnate-leaved palms in Puerto Rico, *Acrocomia media* and *Aiphanes acanthophylla*. The latter is rare in the forests of the subtropical moist zone, on the 'haystack hills' in the north of the island. It is easily distinguished, not only by its spines, but also by the truncated, jagged ends of the leaflets. The only other spiny palm, *Acrocomia media*, the "palma de corozo," is common at lower elevations in forests and fields. This species has deciduous spines on its trunk, and is found only in Puerto Rico and one of the Virgin Islands, St. Thomas.

In the hills between San Germán and Lajas, *Goussia attenuata* is frequently seen. It grows on ridges and sticks out above the surrounding vegetation, the slender leaning trunk supporting a small head of pinnate leaves. Normally only isolated specimens are seen, and in contrast to palms like *Roystonea*, it disappears when its habitat is disturbed. This palm is



6. *Acrocomia media* is common in lower elevations in forests and fields.

endemic to Puerto Rico, and although not common it has two strongholds, not only in the San German hills but also in the northeast of the island near Manati. The roots of this species are interesting. They are visible above ground and are covered with spirally arranged spine-like projections. There is only one other species, *Gaussia princeps* from Cuba. Like *Thrinax* and *Coccothrinax*, *Gaussia* only grows on calcareous soils, although it is in no way related to these two coryphoid genera, being a member of the chamaedoreoid group of palms.

In the wet forests of the Sierra de Luquillo and the Cordillera Central *Pres-toea montana* (syn. *Euterpe globosa*) is abundant, at altitudes of 300 to 1,000 m. There have been numerous ecological studies in these forests and much is known about the ecology of this palm, the "palma de sierra." It often occurs in dense strands, known as palm brakes, in very wet soil of steep slopes. Some ecologists believe that this palm is a secondary successional

species, invading areas that have been disturbed naturally, either by landslides or hurricanes. Others think that *P. montana* is a typical climax species, and its dominance in steep wet areas is a result of its ability to anchor itself by means of numerous above-ground roots. Set against both these theories is the fact that this palm is very slow growing, only 20–30 cm per year, and despite its tremendous reproductive ability (5,000 seeds a year per mature tree) only one in a million seeds reaches maturity (Bannister 1970). There are also some interesting associations between this species and other animals and plants. The seeds are eaten by scolytid beetles (Janzen 1972) and the flowers are bee-pollinated. The trunks and leaves of the palms are often covered with epiphytes, from lichens, mosses and liverworts, to ferns and flowering plants, such as bromeliads and orchids.

Certainly the rarest Puerto Rican palm is the endemic *Calyptronoma rivalis*. It was discovered in 1901 by O. F. Cook, beside a small stream between Lares and San Sebastian. It is still there today, and has recently been found in a second locality nearby. The original site contains about 30 mature specimens, and since they are all on private land they are in a perilous position. Part of one colony has recently been destroyed, but it is to be hoped that the high level of interest by local botanists will ensure their survival. There are other closely related species of *Calyptronoma* in the Caribbean, and three closely related genera, *Calyptrogyne* in Central America and *Pholidostachys* and *Welfia* in both Central America and northern South America. They are all allied to the large genus *Geonoma*.

The tenth and last Puerto Rican palm has least claim to be included in the flora. One single specimen of *Pseudophoenix sargentii* has been recorded from Mona Island, 35 miles west of the main island. This represents the eastern limit of this species which is found from Hispaniola to Mexico.

Acknowledgments

My thanks to the many botanists in Puerto Rico who were so helpful to me on a recent visit to the island.

LITERATURE CITED

BANNISTER, B. A. 1970. Ecological life cycle of *Euterpe globosa* Gaertn., p. B-299-314. In H. T. Odum and R. F. Pigeon (Eds.). A tropical

rainforest. U.S. Atomic Energy Comm., Washington, D.C.

JANZEN, D. H. 1972. Association of a rainforest palm and seed-eating beetles in Puerto Rico. *Ecology* 53: 258-261.

LITTLE, E. L. AND WADSWORTH, R. H. 1964. Common Trees of Puerto Rico and the Virgin Islands. Agriculture Handbook No. 249. USDA Forest Service.

LITTLE, E. L., WOODBURY, R. O. AND WADSWORTH, F. H. 1974. Trees of Puerto Rico and the Virgin Islands. 2nd Vol. Agricultural Handbook No. 449. USDA Forest Service.

LETTERS

Dear Editors and Contributing Writers,

This is a thank-you note to all the people who contribute their efforts toward making *Principes* the outstanding publication that it is. A prime example in point is the January 1984 issue. That issue has something for everyone.

Who could ask for anything more than to discover, or in this case, rediscover a palm like *Livistona exigua* in Borneo? How about the possibility of an indoor *Livistona* that looks like a *Licuala*? Thanks to P. R. Morgan and John Dransfield!

How about an article on the thirteen things that could possibly be wrong with your palm? You never knew whom to ask before, because no one had done the research. Now, thanks to T. K. Broschat and the University of Florida Agricultural Research and Education Center, we know who and where to ask. Thanks for making our palms green and healthy!

Next, we boarded our worldwide traveling armchair to Nepal. Here, Melvin and Phyllis Sneed guided us past Mt. Everest, then from Kathmandu, via elephant, land-rover, and foot to see strange *Phoenix* palms, rare one-horned rhinos, and the Royal Botanical Gardens. Who wouldn't love to go along with the Sneeds? Thanks for sharing your adventures with us!

Perhaps the ultimate plant of all, the Coconut Palm, is discussed in the next article by Pablo Guzman-Rivas. The writer talks of other palms of Mexico and the

Philippines also. Who could ever tire of reading more about the history and versatility of *Cocos nucifera*? Gracias amigo!

And then, Nancy G. Dengler and Ronald E. Dengler explore the microscopic formation of plications in two palm leaves. Everyone should have a scanning electron microscope. Thanks to these two people for taking us into the microscopic world where the folding begins to occur. Without the scientist and those who ask why, we would all still be in the Dark Ages.

And then there is the news. Of course, Florida and California have it all, but can you imagine Texas, the almost largest state with the smallest chapter, and that new chapter in Ohio, burrrrrrrrr!

And of course, our very own bookstore with the latest and greatest literature about palms. For \$1.25 plus cost, you can build up your palm library.

In summary, we find that within this *Principes*, there is discovery, and/or, rediscovery of palms, how-to/or how not-to grow palms, a travelog, history and use of palms, and last but not least, the nitty-gritty of leaf formation. All of the above plus classified ads, Society news, a bookstore, and a Palmeter. What Dent Smith started and Harold E. Moore worked for (along with countless others) has turned into a marvelous Society with a truly remarkable publication, *Principes*. Thank you all!

Sincerely yours,
ERWIN M. RUHLAND

Where's Joey?

MELVIN W. SNEED

8107 S.W. 72nd Ave., Apt. 113 E, Miami, Florida 33143, U.S.A.

Once upon a time there was a palm, and for more than a century it was named *Teysmannia*. Early on, it was thought to be a genus with a single species; namely, *T. altifrons*. Time has proved it to be one of the world's most beautiful palms, with three other species described. But very few collectors have this palm in their gardens, nor can one find it in most of the world's renowned botanical gardens. Extremely "rare in cultivation" is the cachet attached to it.

The name was changed, perhaps with good reason, as explained by our late editor of *Principes*, Dr. Harold E. Moore, Jr. (see Vol. 5, No. 4, 1961, p. 116). So the palm subsequently has been known as *Johannesteijsmannia*, and the name change (which indeed is a "mouthful") also provided a cover for additional species which have been identified and described by Dr. John Dransfield.

During some of our travels, Phyllis Sneed and I went in quest of this palm, found it, and tried to collect it, but with little success. Seeds, especially viable ones, are extremely hard to find. Except for one of the species, which Dr. John Dransfield discovered up north in Malaysia west of Ipoh, the palm has no trunk. The large, entire, diamond-shaped leaves are stiff and leathery in texture, extending up from the ground on long leaf stalks. *J. altifrons* fruits at ground level, so collectors beware! When brushing away debris from the base of the plant to look for seeds, use a stick rather than hand to avoid contact with a viper. But do not be deterred. Your author never has seen a snake in the forests where

this tree occurs, though no doubt some are there.

As one might surmise, the genus is indigenous largely to Malaysia (Peninsula Malaysia and Sarawak); it also occurs in Sumatra. If one crosses the causeway from Singapore to Malaysia, he will be in Johore; then a relatively short drive on up the East Coast to Mersing will deliver the "Joey" seeker into an area where they are abundant. However don't imagine that seeds will crawl into your open basket. Even here one should desert the highway and follow old loggers' roads back into



1. *Johannesteijsmannia* in the forest, Borneo.



2. Rain will not penetrate or leak through any shelter properly thatched with "Joey's" leaves.




what's left of the forest. That would be the best chance of finding colonies of "Joey," some of which might be in fruit.

Also one can find this palm dispersed in forests in north and central western Malaysia as well as the eastern Malaysian State of Sarawak, Borneo. (See Fig. 1 for a host of them under the forest canopy in Bako National Park, Sarawak.) We observed that "Joey's" leaf is one of the best possible thatches for native structures. One does not have to shred it, mat it, or weave it. Just pluck the magnificent leaf and attach it to whatever framework is to be covered, overlap the leaves slightly,

←

3. *Johannesteijsmannia* in the Botanic Garden, Singapore (Eric Taylor furnishes scale). This may be the first "Joey" palm cultivated in the Garden. At one time, Waterfall Gardens, in Penang, had one growing, but it was lost.



4. Recently, the Singapore Garden has cultivated and planted out in two open areas (small triangular plots) three "Joey" in each.

and one has a waterproofed shelter (Fig. 2).

So we've seen the palm in its habitat. Where is "Joey" elsewhere? In 1978 we admired one specimen planted in the extensive gardens of Palm Society member, Mr. Paul Berli, in Bangkok, but it no longer survived in 1983 when we returned. To our knowledge, only the Singapore Botanical Garden now can boast of having it set out and growing.

Historically, some director got a "Joey" transplanted and established near his house at the far end of the Singapore Garden. Actually, this area was "out-of-bounds" for the multitudes visiting the garden, so hardly anyone saw the palm there. But the plant thrived, and on our second visit to the Garden we found and photographed it (Fig. 3). Since then the Singapore Garden, realizing that it has one of the world's

rarest and most beautiful palms in its own backyard, has cultivated the palm and now has six new plants set out, which we discovered in April, 1984. Three are set out in a shade area adjacent to young *Verschaffeltia splendida*, easily visible from a walkway, and three others are in deepest shade across a walkway from the orchid houses (Figs. 4, 5).

So far we have been experiencing *Johannesteijsmannia altifrons*. But with thanks to John Dransfield's field work, there are other species of this unusual palm to be sought. As recently as April, 1984, we were looking for palms in the Cameron Highlands, some five hours leisurely drive north of Kuala Lumpur, capital of Malaysia. Once there we were not far from Ipoh, in the outstretches of which Dr. Dransfield found the single trunked "Joey."

We wanted to drive on to Ipoh, but our



5. Same as Fig. 4, with your author admiring a specimen.



6. More palms, including *Iguanura*, on the hill (photographed by Eric Taylor).

companion, and long-time Palm Society member in Malaysia, Eric Taylor, said, "Mel, we can't get in and out of Ipoh. The Malaysian Government has some insurgents bottled up there, and we are not able to get into the area where the palm is without special permission from the Government." That squelched the Ipoh end of our trek, but we had fun looking in other directions.

For three days we explored the beauty of Cameron Highlands and collected seed for the Palm Society Seed Bank, most notably of *Livistona speciosa*, *Licuala spinosa* and *Orania sylvicola*. This area is loaded with many species of palms, though we saw no "Joeys." We found *Iguanura wallichiana* (Fig. 6) and collected a few seeds of questionable viability, and the area is rich in species of *Pinanga*.

Back to Kuala Lumpur, Eric drove us south to Seremban, passing plantations of *Metroxylon*, *Elaeis guineensis*, coconuts, mangoes, mangosteens and longsats. We



7. Eric Taylor, Azhari Ahmad and your author pose with leaves of *Licuala longipes* and *Johannesteijsmannia magnifica*.



8. Underside of leaf of *Johannesteijsmannia magnifica*.

saw some of the huge durian trees along the roadside as well as one fruiting *Actinorhysis calapparis*. We passed a cocoa factory where Cadbury chocolate is refined from nearby cocoa plantings. Turning back north and west to the Jelebu area, we arrived at Jeram Toi picnic area by a waterfall with cement steps leading about 200 feet along a tumbling stream, and steep hills above it. Up the hillside and atop were *Johannesteijsmannia magnifica*, *Licuala longipes* and *Pinanga* sp. (Fig. 7).

The front side of this "Joey" leaf was a deep green, pleated and stiff. The reverse side was even more beautiful, silver-tinted, glistening in the sun, with perfect configuration and displaying a natural etching that borders the entire leaf in a zig-zag design (See Fig. 8). This species is the "grey" *Teysmannia* referred to by Dr. T. C. Whitmore in *Palms of Malaya*, p. 111, and only recently named by John Dransfield.

So, where do we go from here? Without doubt, *Johannesteijsmannia* being one of the world's most unusual and beautiful palms is endangered by logging and the shrinking forests of its habitat. In retrospect, the only way this rare palm ever will have a chance of being distributed is through the Palm Society Seed Bank. It behooves all of us (Palm Society members) to think a bit, and mount and support some effort to capture seeds of "Joey" to get it growing someplace outside of Malaysia. It will take a lot of "doing"!

BOOKSTORE

A MANUAL OF THE RATTANS OF THE MALAY PENINSULA (J. Dransfield 1979, 270 pp.)	\$25.00
COCONUT PALM FROND WEAVING (Wm. H. Goodloe 1972, 132 pp.)	3.95
CULTIVATED PALMS OF VENEZUELA (A. Braun 1970, 94 pp. and 95 photographs.)	4.50
FLORA OF PANAMA (Palms) (R. E. Woodson, Jr., R. W. Schery 1943, 122 pp.)	17.00
FLORA OF PERU (Palms) (J. F. MacBride 1960, 97 pp.)	8.00
FLORIDA PALMS, Handbook of (B. McGeachy 1955, 62 pp.)	1.95
HARVEST OF THE PALM (J. J. Fox 1977, 244 pp.)	16.50
INDEX TO PRINCIPES (Vols. 1-20, 1956-1976, H. E. Moore, Jr., 68 pp.)	3.00
MAJOR TRENDS OF EVOLUTION IN PALMS (H. E. Moore, Jr., N. W. Uhl 1982, 69 pp.)	6.00
PALMS (A. Blombery & T. Rodd 1982, 192 pp., 212 colored photographs)	30.00
PALMS IN AUSTRALIA (David Jones 1984, 278 pp., over 200 color photographs)	25.00
PALMS FOR THE HOME AND GARDEN (L. Stewart 1981, 72 pp., some color)	10.95
PALMS OF BRITISH INDIA & CEYLON (Blatter 1926, reprinted in India 1978, 600 pp.)	75.00
PALMS OF THE LESSER ANTILLES (R. W. Read 1979, 48 pp.)	8.00
PALMS OF MALAYA (T. C. Whitmore 1973, 132 pp.)	16.95
PALMS OF SOUTH FLORIDA (G. B. Stevenson 1974, 251 pp.)	7.95
PALMS OF THE WORLD (J. C. McCurrach 1960, 290 pp.)	19.00

In Malaysia, spring of 1984, we made an arrangement with Eric Taylor to correspond with Fairchild Garden, here in Miami, Florida, with a view to a plant exchange, including a seedling of "Joey." Eric had two of them started in his home nursery in Kuala Lumpur. We hope, eventually, that this deal will work out and Fairchild Garden will be a well-deserved beneficiary, with all thanks due to Eric Taylor.

PALM SAGO (K. Ruddle, D. Johnson, P. K. Townsend, J. D. Rees 1978, 190 pp.)	7.50
SECRET OF THE ORIENT DWARF <i>Rhapis excelsa</i> (L. McKamey 1983, 51 pp.)	3.95
SUPPLEMENT TO PALMS OF THE WORLD (A. C. Langlois 1976, 252 pp.)	25.00
THE GENUS <i>PTYCHOSPERMA</i> LABILL. (F. B. Essig 1978, 61 pp.)	5.50
THE INDIGENOUS PALMS OF SURINAME (J. G. W. Boer 1965, Part of Flora, 172 pp.)	42.00
THE MAJOR GROUPS OF PALMS AND THEIR DISTRIBUTION (H. E. Moore, Jr., 1973, 115 pp.)	4.50
THE MINIATURE PALMS OF JAPAN (U. Okita, J. L. Hollenberg 1981, 135 pp.)	19.95
THE PALM FLORA OF NEW GUINEA (F. B. Essig, 1977, 46 pp.)	6.50

PALM PAPERS (Postage Included)

FURTHER INFORMATION ON HARDY PALMS (J. Popenoe 1973, 4 pp.)	1.25
NOTES ON PRITCHARDIA IN HAWAII (D. Hodel 1980, 16 pp.)	2.00
RARE PALMS IN ARGENTINA (reprint from <i>Principes</i> , E. J. Pingitore 1982, 9 pp., 5 beautiful drawings)	2.75
THE HARDEST PALMS (J. Popenoe 1973, 4 pp.)	1.25

The palm books listed above may be ordered at the prices indicated plus \$1.25 extra per book to cover packaging and postage. (California residents please add 6% sales tax.) Foreign checks must be in US dollars and payable on a USA bank. In some countries it is possible to send International Money Orders through the Post Office. Send check payable to The International Palm Society to Pauline Sullivan, 3616 Mound Avenue, Ventura, CA 93003, U.S.A. ALL SALES FINAL.

Vessel Elements in Roots of Young Palms

MONORANJAN GHOSE

Indian Statistical Institute, Calcutta-700 035, India

ABSTRACT

Perforation plate type, end wall slope, and length to width ratio are described in wide vessel elements from roots of seedlings of 17 species of palms. Vessel elements with two different (i.e., "mixed") perforation plate types or end wall slopes occur in many of the specimens. The data contain correlations between the end wall characters and the length to width ratio of vessel elements. There is little resemblance between the sequence of the species according to Moore's (1973) systematic classification of palms, and the sequence according to the length to width ratio of vessel elements. The wide vessel elements from roots of young palms are usually more variable than those from roots of adult palms of the same species.

Data have been published on tracheary elements in the organs of adult palms (Bierhorst and Zamora 1965; Cheadle 1942, 1943*a*, 1943*b*; Klotz 1978*a*, 1978*b*, 1978*c*; Parthasarathy and Klotz 1976; Tomlinson 1961; Tomlinson and Zimmermann 1967). In seedling palms, no comparable work appears to have been done so far. The present study describes perforation plates, end wall slopes, lengths, widths, and length-width ratios of vessel elements in roots of seedling palms. The objectives are (1) to compare the data from young palms with those presented by Tomlinson (1961) and Klotz (1977) from adult palms, (2) to determine the degree to which end wall characteristics are correlated with length to width ratios, and (3) to observe whether the arrangement of species according to length to width ratios of vessel members resembles the systematic sequence of taxa in Moore's (1973) classification of palms.


The form of tracheary elements and the correlations among their features have supplied evidence for hypothetical evolu-

tionary pathways (Carlquist 1975; Dickison 1975). Recently, Aldridge (1978) used a method of statistical correlation between features of tracheary elements in studying evolution within the genus *Sonchus*. The present analysis is based on her hypothesis about the evolution of tracheary elements. She used the length to width ratio of vessel elements as a tool to determine the phylogenetic relationship among species. A higher ratio indicates primitiveness while a lower ratio is more advanced (Aldridge 1978). According to Cheadle (1943*a*), primitive vessel elements have long, extremely oblique or very oblique end walls in which the perforation plates have many bars that are closely spaced while advanced vessel elements have slightly oblique to transverse end walls on which the bars are few or absent.

Materials and Methods

Small pieces of mature roots of average diameter were studied in seedlings of 17 species of palms (Table 1). Most of the seedlings were raised at the Crop Garden of the Indian Statistical Institute, Calcutta, from seeds collected from the Indian Botanic Garden, Howrah. The seeds of *Veitchia* and *Phoenix rupicola* were collected in Australia. Seeds were sown at different times according to their availability. The ages of the seedlings ranged from 8 to 30 months.

One piece of a root was sampled for each species. The roots were cut into small pieces and fixed in formalin-acetic acid-alcohol (FAA) for 48 hours or more. They were then boiled in 10% potassium



1. Camera lucida drawings of wide vessel elements of roots. Lateral-wall pits are represented as short, horizontal lines in A_2 and B_1 . Magnification is $107\times$ in A_2 and B_1 , $43\times$ in all the others. A_1 , A_2 . *Areca catechu*. A_1 . Simple perforation plates, transverse end walls. A_2 . Mixed perforation plates and end wall slopes. B_1 , B_2 . *Chrysalidocarpus lutescens*. B_1 . Details of scalariform perforation plate. B_2 . Simple perforation plates, transverse end walls. C_1 , C_2 . *Livistona rotundifolia*. C_1 . Simple perforation plates, transverse end walls. C_2 . Mixed end wall characters. D . *Arenga pinnata*. Simple perforation plates, transverse end walls. E_1 , E_2 . *Borassus flabellifer*. E_1 . Simple perforation plates, end wall slopes mixed. E_2 . Simple perforation plates, transverse end walls. F . *Cocos nucifera*. Simple perforation plates, transverse end walls. G_1 , G_2 . *Areca triandra*. G_1 . Simple perforation plates, transverse end walls. G_2 . Mixed end wall characters. H_1 , H_2 . *Veitchia merrillii*. H_1 . Mixed end wall characters. H_2 . Simple perforation plates, transverse end walls. I . *Phoenix rupicola*. Simple perforation plates, mixed end wall slopes. J_1 , J_2 . *P. reclinata*. J_1 . Mixed end wall characters. J_2 . Simple perforation plates, transverse end walls. K . *Salacca zalacca*, mixed end wall slopes, simple perforation plates. L . *Elaeis guineensis*, mixed end wall characters. M_1 , M_2 . *Hyphaene dichotoma*. M_1 . Mixed end wall characters. M_2 . Simple perforation plates and transverse end walls.

hydroxide (10 minutes), washed in water, soaked in 25% chromic acid (15 minutes), and washed again. The macerated material was teased apart and mounted on slides in phenol-glycerine. Some specimens fixed in FAA were sectioned transversely with a microtome and stained with safranin and fast-green. Camera lucida drawings and photomicrographs were made.

An ocular micrometer in a compound microscope was used to measure the length and width of 10 intact wide (or sometimes semi-wide) vessel elements in each species (Fig. 3). Slopes of end walls and types of

perforation plates were also recorded. The mean and standard error of the measurements were computed. Spearman's rank correlation coefficient was employed to demonstrate the correlation between the end wall characteristics and length to width ratios.

Categories of Perforation Plates and End Wall Slopes

The terminology for perforation plates (except "mixed") is taken from Esau (1958), and the vessel elements are cat-

Table 1. Perforation plate types, end wall slopes, and dimensions of wide metaxylem vessel elements in roots of young palms.¹

Major groups and species	% of vessel elements with perforation plates		% of vessel elements with end walls			Dimension of vessel elements						
	Both scalariform	Mixed	Both simple	Both oblique	Both slightly oblique	Mixed	Both transverse	Length (mm)		Width (mm)		Length : width rate
								Mean ± std. error				
Coryphoid												
<i> Livistona rotundifolia</i> (Lam.) Mart. (10)	—	20	80	—	—	30	70	0.62 ± 0.04	0.062 ± 0.006	11.9 ± 2.1		
Phoenicoid												
<i> Phoenix reclinata</i> Jacq. (8)	* 10	20	70	10	10	50	30	1.12 ± 0.07	0.042 ± 0.005	28.5 ± 3.8		
<i> P. rupicola</i> Anders. (10)	10	10	80	10	40	—	50	2.05 ± 0.13	0.090 ± 0.004	23.5 ± 2.0		
<i> P. sybestrus</i> (L.) Roxb. (16)	—	—	100	—	—	—	100	1.84 ± 0.10	0.091 ± 0.002	20.5 ± 1.1		
<i> P. pusilla</i> Gaertn. (16)	—	20	80	10	40	10	40	1.07 ± 0.08	0.079 ± 0.010	14.9 ± 1.7		
Borassoid												
<i> Borassus flabellifer</i> L. (8)	—	10	90	20	10	30	40	1.40 ± 0.15	0.064 ± 0.002	22.6 ± 3.0		
<i> Hyphaene dichotoma</i> (White) Furtado (15)	10	10	80	10	—	20	70	2.16 ± 0.16	0.104 ± 0.009	23.3 ± 4.0		
Lepidocarpyoid												
<i> Salacca zalacca</i> (Gaertn.) Yoss (12)	—	20	80	—	—	20	80	1.52 ± 0.11	0.090 ± 0.005	18.8 ± 3.6		
Caryotoid												
<i> Arenga pinnata</i> (Wurmb.) Merr. (30)	—	—	100	—	—	—	100	1.02 ± 0.06	0.091 ± 0.009	11.3 ± 0.9		
<i> Caryota urens</i> L. (12)	—	—	100	—	—	—	100	2.39 ± 0.24	0.189 ± 0.023	15.0 ± 3.4		
Arecoïd												
<i> Areca catechu</i> L. (24)	10	10	80	—	10	10	80	0.88 ± 0.10	0.077 ± 0.005	12.3 ± 2.1		
<i> A. triandra</i> Roxb. (19)	—	40	60	—	—	40	60	1.02 ± 0.12	0.068 ± 0.004	16.8 ± 3.9		
<i> Chrysalidocarpus lutescens</i> H. A. Wendl. (16)	—	10	90	10	—	—	90	1.90 ± 0.16	0.084 ± 0.003	22.9 ± 2.2		
<i> Roystonea regia</i> (H.B.K.) Cook. (11)	—	—	100	10	—	—	90	1.18 ± 0.09	0.111 ± 0.004	10.7 ± 0.9		
<i> Veitchia merrillii</i> (Becc.) Moore (10)	—	20	80	—	—	40	60	2.15 ± 0.14	0.064 ± 0.003	33.7 ± 2.2		
Cocosoid												
<i> Cocos nucifera</i> L. (12)	—	—	100	—	—	—	100	1.86 ± 0.18	0.200 ± 0.017	10.4 ± 1.8		
<i> Elaeis guineensis</i> Jacq. (12) ²	30	40	30	15	40	30	15	1.03 ± 0.07	0.043 ± 0.003	24.7 ± 1.8		
Total of species	5	12	17	8	6	10	17	—	—	—		

¹ The classification is that of Moore (1973). The age of the plant in months is given in parentheses after the binomial.² Only 7 vessel elements were examined in *Elaeis guineensis*.

egorized as follows: both perforation plates scalariform; "mixed" perforation plates, i.e., one plate simple and the other scalariform or reticulate (Fig. 1A₂, C₂, G₂, H₁, J₁, L, M₁); both perforation plates simple (Fig. 1A₁, B₂, C₁, D, etc.). Similarly, Cheadle's (1943a) classification of end wall slopes is employed, and the vessel elements are categorized as follows: both end walls oblique; both end walls slightly oblique; "mixed" end wall slope, i.e., one end wall transverse, and the other either oblique or slightly oblique (Fig. 1A₂, C₂, E₁, G₂, H₁, I, L, etc.); both end walls transverse (Fig. 1A₁, B₂, C₁, D, E₂, F, G₁, etc.).

Observations

Simple perforation plates and transverse slopes are the most frequent end wall characters in the vessel elements examined in this survey. Mixed perforation plates and mixed end wall slopes are the next most frequent. Table 1 shows that 17 species have vessel elements with two simple perforation plates; 12 species, mixed perforation plates; and five species, two scalariform perforation plates. *Elaeis* shows the highest percentage (30%) of scalariform perforation plates. The highest percentages of vessel elements with mixed perforation plates occur in *Elaeis* (40%) and *Areca triandra* (40%). The percentage of vessel elements with two simple perforation plates is highest in all of the species except *Elaeis*.

Vessel elements with both end walls oblique occur in eight species, both slightly oblique in six species, and both transverse in all the species. Vessel elements with mixed slopes occur in 10 species. The highest frequencies of vessel elements with both end walls slightly oblique occur in *Elaeis*, *Phoenix rupicola* and *P. pusilla*. *Areca triandra* (40%), *Phoenix reclinata* (50%), and *Veitchia* (40%) show the highest percentages of vessel elements with mixed slopes. The percentage of vessel

elements with two transverse end walls is highest in all of the species except *Elaeis* and *Phoenix reclinata*. The highest numbers of vessel elements with both perforation plates simple and both end walls transverse occur in *Arenga*, *Caryota*, *Cocos*, *Phoenix sylvestris* and *Roystonea*.

The dimensions and length to width ratios of the vessel elements are also given in Table 1. The longest is found in *Caryota*, the shortest in *Livistona*. On the other hand, the widest is in *Cocos* and the narrowest are in *Elaeis* and *Phoenix reclinata*. The highest mean ratio occurs in *Veitchia* and the lowest in *Cocos*.


Spearman's coefficient of rank correlation (Steel and Torrie 1960) was employed to demonstrate correlations between the length to width ratio and the end wall characters. The data for these calculations are displayed graphically in Figure 2. The 17 species were ranked in increasing order of the average length to width ratio, and in a specialization index (adapted from Klotz 1978) based on end wall slopes (Fig. 2B):

Specialization index of end wall slopes \times 100 = 1 (% of both end walls transverse) + 2 (% of mixed end walls) + 3 (% of both end walls slightly oblique) + 4 (% of both end walls oblique).

The Spearman's coefficient of rank correlation between these two features is 0.650, which is significant at the 1% level. Similarly, the following specialization index was assigned for perforation plates (Fig. 2A):

Specialization index of perforation plates \times 100 = 1 (% of both plates simple) + 2 (% of mixed plates) + 3 (% of both plates scalariform).

The Spearman's coefficient of rank correlation (with length to width ratio) is 0.525, which is significant at the 5% level. In summary, the species with the lower length to width ratios tend to possess the



2. Correlation of length to width ratios with end wall features. A. Specialization indices of perforation plates plotted against length to width ratios. B. Specialization indices of end wall slopes plotted against length to width ratios.


higher percentages of simple perforation plates and transverse end walls. In contrast, the higher length to width ratios tend to occur in the species with the higher percentages of mixed end wall characters. A few species are exceptions to these trends (e.g., *Livistona*, *Phoenix sylvestris*, and *P. pusilla*).

Table 2 shows the differences in wide vessel element dimensions and end wall characteristics between young and adult palms. The young plants possess various types of perforation plates, the simple type predominating, whereas adult plants possess only the simple type. Again, various types of end wall slopes occur in young plants, but only transverse or sometimes slightly oblique in adults. The range of lengths of vessel elements is generally larger in the young plants than in the adults. The upper limit of length of vessel

elements in young plants exceeds that in adult plants except in *Elaeis* and *Phoenix*. In contrast, the widths of the vessel elements are mostly greater in the adult plants.

Discussion

The data in the present study differ from the comparable observations by Tomlinson (1961) and Klotz (1977) in adult palms of the same species. The comparisons between young and adult palms reveal a general trend of shortening and widening of the wide vessel elements in the roots. Also, the end wall characters of these vessel elements are more variable in the young palms than in the adult palms. Thus, the pattern of differentiation of these cells apparently undergoes a transformation during the development of an individ-



3. Transverse section of root of *Phoenix pusilla*. $\times 100$. e, endodermis; n, narrow tracheary elements; ph, phloem; sw, semi-wide vessels of metaxylem; w, wide vessels of metaxylem.

ual palm from the seedling stage to the adult. In the above features, the wide vessel elements from seedling roots resemble the narrow, early-maturing vessel elements from roots of adult palms (described by Cheadle 1942; Klotz 1977, 1978a; Parthasarathy and Klotz 1976).

Tomlinson (1961), Cheadle (1942, 1943a, 1973b) and Klotz (1978a) did not mention any mixed end wall characters in vessel elements from the species of adult palms that they studied. Apparently, such vessel elements either are absent in adult palms or have not been reported by the previous authors. However, Tomlinson and Zimmermann (1976) depicted mixed end wall characters in vessel elements from the lower part of the stem of *Sabal palmetto*; and Klotz (pers. comm.) observed but did not describe such vessel elements in the vegetative organs of various palms.

Klotz (1977, 1978a) observed reticulate and/or scalariform-reticulate perforation plates in wide vessel elements from

Table 2. Comparison of wide vessel elements from roots of young and adult palms.

Species	Author's data (young plants)		Tomlinson's (1961) data (adult plants)		Klotz's (1977) data (adult plants)	
	Dimensions (μm)	End wall*	Dimensions (μm)	End wall*	Dimensions (μm)	End wall*
<i>Phoenix reclinata</i>	900-1,500 \times 30-60	sc, si/o, s, t	—	—	1,000-1,900 \times 140-260	si/t
<i>Salacca zalacca</i> (<i>S. edulis</i>)	1,000-2,200 \times 40-100	sc, si/s, t	1,000-1,400 \times 180-260	si/t	—	—
<i>Areca catechu</i>	600-1,600 \times 60-100	sc, si/s, t	840-1,210 \times 115-190	si/t	—	—
<i>Chrysalidocarpus lutescens</i>	1,400-2,800 \times 60-90	sc, si/o, t	1,400-1,800 \times 95-210	si/t	—	—
<i>Cocos nucifera</i>	900-3,100 \times 130-270	si/t	1,740-2,740 \times 200-315	si/t	1,100-2,500 \times 200-230	si/s, t
<i>Elaeis guineensis</i>	700-1,300 \times 30-50	sc, si/o, s, t	1,210-1,500 \times 270-340	si/t	—	—

* End wall characters: sc = scalariform perforation plates; si = simple perforation plates; si = slightly oblique end walls; s = oblique end walls; o = oblique end walls; t = transverse end wall slopes.

roots of some species of adult palms. Such perforation plates were not observed in the present study.

Vessel elements from roots of young palms exhibit positive correlations between the length to width ratio of the cell and the degree of primitiveness of the perforation plates. This finding supports the conclusions of Aldridge (1978) and Cheadle (1943a) that were summarized in the introduction of this paper.

The sequence of mean length to width ratios in Table 1 can be compared with the sequence of species according to Moore's (1973) classification of palms. The lowest ratios occur in species of cocosoid, arecoid, and lepidocaryoid groups, while the highest ratios occur in species of arecoid and coryphoid groups. There is little resemblance between Moore's (1973) systematic sequence of the evolutionary lines and major groups and the sequence of species based on length to width ratio of vessel elements. For example, *Livistona* and *Phoenix pusilla* show advanced ratios although they belong to primitive coryphoid and phoenicoid groups, respectively. Similarly, *Veitchia* and *Chrysalidocarpus* of the advanced arecoid group and *Elaeis* of the advanced cocosoid group show a high ratio, which is a primitive character.

There is considerable variation among the species of individual major groups in characters of the vessel elements. For example, within the cocosoid major group, *Cocos nucifera* shows relatively advanced characters and *Elaeis guineensis*, more primitive characters. Similarly, within the arecoid major group, more advanced xylem occurs in *Roystonea regia* and more primitive in *Veitchia merrillii*. In phoenicoid major group, more advanced xylem occurs in *Phoenix pusilla* and more primitive in *P. reclinata*. Such differences among the members of individual major groups may be explained by Klotz's (1978a) conclusion that "evolution in the xylem appears to have progressed inde-

pendently in the various taxonomic groups of palms."

Acknowledgments

I am grateful to Professor T. A. Davis, former Head, Crop Science Unit, Indian Statistical Institute, for his guidance during the course of this investigation; to Professor B. M. Johri, Delhi University, for his valuable comments on the article; and to Dr. Larry H. Klotz, Shippensburg University, Shippensburg, Pennsylvania, for his comments and scrutiny for improvement of this article.

I thank Mr. S. K. De, Artist and Dr. A. Dasgupta, Senior Research Fellow, Indian Statistical Institute, for drawings and statistical calculations, respectively.

LITERATURE CITED

- ALDRIDGE, A. E. 1978. Anatomy and evolution in the Macaronesian *Sonchus* subgenus *Dendrosonchus* (Compositae Lactuceae). Bot. J. Linn. Soc. 76: 249-285.
- BIERHORST, D. W. AND P. M. ZAMORA. 1965. Primary xylem elements and element associations in angiosperms. Amer. J. Bot. 52: 657-710.
- CARLQUIST, S. 1975. Ecological Strategies of Xylem Evolution. University of California Press, Berkeley.
- CHEADLE, V. I. 1942. The occurrence and types of vessels in the various organs of the plant in the Monocotyledoneae. Amer. J. Bot. 29: 441-450.
- . 1943a. The origin and certain trends of specialization of the vessel in the Monocotyledoneae. Amer. J. Bot. 30: 11-17.
- . 1943b. Vessel specialization in the late metaxylem of the various organs in the Monocotyledoneae. Amer. J. Bot. 30: 484-490.
- DICKISON, W. C. 1975. The bases of angiosperm phylogeny: Vegetative anatomy. Missouri Bot. Gard. 62: 590-620.
- ESAU, K. 1958. Plant anatomy. John Wiley & Sons, New York.
- KLOTZ, L. H. 1977. A systematic survey of the morphology of tracheary elements in palms. Unpubl. Ph.D. Thesis, Cornell University, Ithaca, New York.
- . 1978a. Form of the perforation plates in the wide vessels of metaxylem in palms. J. Arnold Arbor. 59: 105-128.
- . 1978b. Observations on diameters of vessels in stems of palms. Principes 22: 99-106.

- . 1978c. The number of wide vessels in petiolar vascular bundles of palms: an anatomical feature of systematic significance. *Principes* 22: 64–69.
- MOORE, H. E., JR. 1973. The major groups of palms and their distribution. *Gentes Herb.* 11: 27–141.
- PARTHASARATHY, M. V. AND L. H. KLOTZ. 1976. Palm "Wood." I. Anatomical aspects. *Wood Sci. Tech.* 10: 215–229.
- STEEL, R. G. D. AND J. H. TORRIE. 1960. Principles and procedures of statistics. McGraw-Hill Book Company, Inc., New York.
- TOMLINSON, P. B. 1961. Anatomy of the monocotyledons. Vol. II. *Palmae*. Oxford University Press, London.
- AND M. H. ZIMMERMANN. 1967. The "Wood" of monocotyledons. *Int. Assoc. Wood Anat. Bull.* 1967(2): 4–24.

PALM LITERATURE

A BIBLIOGRAPHY OF GRADUATE THESES ON PALMS, PART II

An initial compilation of 101 titles appeared in a previous issue of this journal (Vol. 27(2), 1983, pp. 85–88). The results of continued research into this primary source of information on palms, following the same criteria, now justifies a second installment.

This listing consists of 57 additional doctoral and master's theses. More than one-half of the total were completed in three countries: United States (14), The Philippines (12) and France (10). As before, the major economic palms dominate the subject matter: coconut (23), oil palm (11) and date palm (8). Although most of the studies were completed within the past 15 years, a few older titles were found, such as the dissertation by Gatín dated 1906. Once again I must acknowledge the assistance of numerous individuals who provided me with thesis titles.

Doctoral Theses

- AMMAR, S. 1978. La Culture de tissus de plantes issues de graines appliquée à la multiplication végétative du palmier dattier (*Phoenix dactylifera* L.). University of Tunis, Tunisia.
- ANDERSON, A. B. 1983. The biology of *Orbignya martiana* (Palmae), a tropical dry forest dominant in Brazil. University of Florida, Gainesville.
- ANNET, E. 1921. Contribution à l'étude du palmier à huile. University of Paris, France.
- BALICK, M. J. 1980. The biology and economics of the *Oenocarpus-Jessenia* (Palmae) complex. Harvard University, Cambridge.
- BASU, S. K. 1981. Studies on the palms of Indian Botanic Garden, Calcutta and its vicinity. University of Calcutta, India.
- CARCALLAS, C. D. 1981. Effects of fertilization on yield and yield components of peanut intercrop and on leaf nutrient concentrations of dwarf coconut cultivars. University of the Philippines, Los Baños.
- DACHLIAN, C. P. 1977. Coryphoid palms from the Lower and Middle Eocene of southeastern North America. University of Texas, Austin.
- DESASSIS, A. 1961. Sur les modalités de formation des matières grasses dans le fruit d'*Elaeis guineensis* Jacq. University of Paris, France.
- DRIRA, N. 1981. Multiplication végétative et micropropagation du palmier dattier (*Phoenix dactylifera* L.) à partir d'organes prélevés sur la phase adulte, cultivés "in vitro." University of Tunis, Tunisia.
- FEATHER, T. V. 1982. Occurrence, etiology and control of wilt and dieback of *Phoenix canariensis* in California. University of California, Riverside.
- GATIN, C. L. 1906. Recherches anatomiques et chimiques sur la germination des palmiers. University of Paris, France.
- GHOSE, M. 1982. Anatomy of the vegetative organs of young palms. University of Calcutta, India.
- HAIBU, T. K. 1981. La Culture *in vitro* des tissus de cocotier (*Cocos nucifera* L.). University of Paris, France.
- MYERS, R. L. 1981. The ecology of low diversity palm swamps near Tortuguero, Costa Rica. University of Florida, Gainesville.
- NAIR, S. 1975. Studies on the microflora of the root region of plantation crops, coconut and cocoa. Indian Agricultural Research Institute, Delhi.
- NGUYEA, T. 1981. Anther culture in coconut (*Cocos nucifera* L.). University of the Philippines, Los Baños.
- NJIKE, F. 1976. Les Conditions de développement de la culture du palmier à huile et du cocotier dans le sud de la Côte-d'Ivoire; la problématique de la planification. University of Paris, France.
- NYBERG, A. J. 1968. The Philippine coconut industry. Cornell University, Ithaca.

- OSMAN, A. M. A. 1974. Xenia and metaxenia studies in the date palm, *Phoenix dactylifera* L. University of California, Riverside.
- PILLET-SCHWARTZ, A. M. 1973. Capitalisme d'état et développement rural en Côte-d'Ivoire, la société pour le développement et l'exploitation du palmier à huile en pays ébrié. University of Paris, France.
- POSAS, O. B. 1981. Grazing versus cut-and-carry trials on goats under coconuts. University of the Philippines, Los Baños.
- PROFIZI, J. P. 1983. Contribution à l'étude des palmiers *Raphia* du sud-Bénin—botanique, écologie, ethnobotanique. University of Montpellier, France.
- QUANSAH, S. T. 1954. A chemical investigation of mineral levels in coconut water in relation to soil, season and yield. Imperial College of Tropical Agriculture, Trinidad.
- RAJARATNAM, J. A. 1971. Boron nutrition of the oil palm. University of Singapore, Singapore.
- REYNOLDS, J. F. 1978. Morphogenesis of palms in vitro. (*Howeia*, *Chamaedorea*, *Phoenix*). University of California, Riverside.
- SAADI, M. 1979. Contribution à la lutte contre le bayoud, fusariose vasculaire du palmier dattier. University of Dijon, France.
- TJEEGA, P. 1974. Les Types d'exploitation du palmier à huile dans la région d'Eseka, Cameroun. University of Paris, France.
- Master's Theses**
- AL-MADENI, M. A. 1983. Study of various parameters involved in palm tissue culture with special emphasis on date palm (*Phoenix dactylifera* L.). California State University, Chico.
- AMARAL FILHO, J. 1983. A economia política do babaçu: um estudo da organização da estrato-indústria do babaçu no Maranhão e suas tendências. Federal University of Pernambuco, Recife, Brazil.
- BEKR, A. J. 1931. Development of the embryo of the date palm. University of California, Berkeley.
- BRIONES, J. B. 1975. A feasibility study on cattle raising in a coconut plantation. Asian Institute of Management, Philippines.
- CHAN, K. W. 1972. The effects of plant growth substances on fruit induction, development and fruit abscission in the oil palm (*Elaeis guineensis* Jacq.). University of Malaya, Kuala Lumpur.
- COOK, T. C. 1959. The distribution of certain species of exotic eucalyptuses, palms and bougainvilleas in western, central and southern Texas. (*Washingtonia*, *Sabal*, *Phoenix*). University of Texas, Austin.
- DA CUNHA, B. 1979. Industrialização integral do côco babaçu: efeitos sobre a renda e o emprego dos pequenos produtores no estado do Maranhão. Federal University of Viçosa, Brazil.
- DAVID, J. R. 1973. A study of the fruit development in open pollinated within-spadix and artificial pollination of coconut, *Cocos nucifera* Linn. Araneta University, Philippines.
- DAVIS, A. M. T. 1942. A study of Boscaje de la Palma in Cameron County, Texas, and of *Sabal texana*. University of Texas, Austin.
- DOLAR, S. G. 1961. The botany of coconut (*Cocos nucifera* L.), the environmental factors and cultural practices affecting its production. Cornell University, Ithaca.
- FERNANDO, E. S. 1981. A revision of the genus *Nenga* (Palmae: Arecoideae). University of Reading, England.
- FERNANDO, N. T. 1972. The coconut oil export industry. Asian Institute of Management, Philippines.
- FOONG, S. F. 1980. Weather-based model for estimating oil palm fruit yield. University of Guelph, Ontario, Canada.
- EL JARRAH, A. 1977. Physico-chemical and histological dynamic changes in khadrawi date fruit. Baghdad University, Iraq.
- LUNTINGAN, H. T. 1981. Nut components of eight coconut cultivars in relation to climatic factors. University of the Philippines, Los Baños.
- MEDEIROS-COSTA, J. T. DE. 1982. As palmeiras (Palmae) nativas em Pernambuco, Brasil. Federal Rural University of Pernambuco, Recife.
- MENDIO LA VARGAS, P. A. 1970. La influencia del frío en la germinación del cocotero. National Agrarian University, Lima, Peru.
- ONG, Y. W. 1973. Some observations on *Ganoderma boninense* Pat., the cause of stem rot on oil palm (*Elaeis guineensis* Jacq.) in West Malaysia. University of Malaya, Kuala Lumpur.
- PATHMARAJ, G. R. 1975. The feasibility of manufacturing coconut coir yarn in Sri Lanka for export. Asian Institute of Management, Philippines.
- PILLAI, V. S. K. 1969. Some observations on *Curvularia eragrostoides* (Henn.) Meyer., the cause of seedling blight of oil palm (*Elaeis guineensis* Jacq.). University of Malaya, Kuala Lumpur.
- SANKAT, C. K. 1975. Development of a system for the mechanical extraction of the kernel from the shell of the coconut. University of the West Indies, Trinidad.
- SANTOS, Z. G. DOS. 1976. Avaliação da fertilidade dos solos da região produtora de côco do estado de Sergipe através da técnica de microparcelas. University of São Paulo, Piracicaba, Brazil.
- SEVIDAL, L. C. 1975. Cultural patterns associated with the coconut. University of the Philippines, Quezon City.
- SINADINOS B. B. 1947. The geographical aspects of coconuts. University of Illinois, Urbana.
- SOBRAL, J. DO P. 1976. Composição química e

- digestibilidade do farelo de côco. University of São Paulo, Piracicaba, Brazil.
- SY-QUIA, L. M., JR. 1972. The market for coconut cream milk, dry cereal baby food and coconut flour. Asian Institute of Management, Philippines.
- TARIGANS, D. D. 1982. Effects of foliar spray of urea on the growth of coconut (*Cocos nucifera*) seedlings. University of the Philippines, Los Baños.
- VALLADE, J. 1965. Recherches morphologiques et cytologiques sur l'embryon d'*Elaeis guineensis* Jacq. quiescent et en cours de germination. University of Dijon, France.
- ZAID, A. 1981. Rapid propagation of the date palm through tissue culture. Institute of Agronomy and Veterinary Medicine, Rabat, Morocco.
- ZARA, D. L. 1981. Tuber production in sweet potato (*Ipomoea batata* (L.) Lan.) intercropped with coconuts: effect of K fertilization and vine pruning. University of the Philippines, Los Baños.

DENNIS JOHNSON

A TOUR AND A COLLECTING TRIP

ROYAL HORTICULTURE SOCIETY TOUR TO THE SEYCHELLES, 12-17 April 1985, led by John Dransfield. For further details contact: ABERCROMBIE AND KENT TRAVEL, 42 Sloan St., London SW1X 9LU, England.

COLLECTING TRIP TO PANAMA. I am planning a trip to Panama, February 1985 to collect PALMS and ORCHIDS. Anyone interested contact JOHN O. WILD, 1123 15th Ave. S.E., Minneapolis, Minn. 51414. (606) 331-7221. Limited to five people.

Principes, 28(4), 1984, pp. 189-190

INDEX

- Acrocomia 21, 23; media 170, 1971; mexicana 21
 Actinorhynchus calapparia 84, 104
 Aiphanes 55; acanthophylla 170; caryotifolia 59; elegans 77;
 pachyclada 62
 Ancistrophyllum 161
 Anderson, Cynthia, as coauthor 55
 Archontophoenix 59, 92, 95, 99, 100
 Areca 21, 131; caliso 20; catechu 16, 20, 84, 104, 119,
 180, 181, 184, 185; ipot 20; triandra 180, 181, 182
 Arecastrum 49, 92, 95, 144; romanzoffianum 90, 127
 Arenga 21, 22, 182; ambong 20; australasica 100; pinnata
 20, 23, 180, 181; porphyrocarpa 103; tremula 20, 23
 Asterogyne 131
 Astrocarum 128; mumbaca 73, 76, 77; standleyanum 128
 Attalea 29; cohune 19; racemosa 75, 76, 77
 Bactris 21, 107, 108, 114, 115, 116, 117, 118, 125, 131;
 gasipaes 59, 105, 106, 107, 108, 109, 110, 112, 113,
 128; granatensis 74, 77; guineensis 78, 117; longiseta
 117; macana 77, 78; major 116, 117; piritu 78;
 porschiana 107, 108, 109, 111, 112, 114, 115, 116;
 setulosa 78, 79; simplicifrons 79, 80; wendlandiana 117
 Basselinia 49, 131
 Beach, James H.
 The reproductive biology of the peach or "Pejibaye" palm
 (*Bactris gasipaes*) and a wild congener (*B. porschiana*)
 in the Atlantic lowlands of Costa Rica 107
 Biennial 1984, Notice of 30, 144
 Bismarckia 49, 84, 103; nobilis 19, 72, 104, 167
 Blomberg, Alec and Rodd, Tony
 Palms (reviewed), 102
 Bookstore 51, 101, 145, 178
 Borassus aethiopicum 155; flabellifer 180, 181
 Brahea 21, 95, 96, 97; armata 19; brandegeei 19; dulcis
 21; nitida 19
 Braun, August
 More Venezuelan palms 73
 Broschat, T. K.
 Nutrient deficiency symptoms in five species of palms grown
 as foliage plants 6
 Broschat, T. K., and Donselman, Henry M.
 Root regeneration in transplanted palms 90 *
 Butia 95
 Butia-Arecastrum hybrid 49
 Calamus 20, 21, 22, 99, 100; deeratus 161
 Calyptrogyne 131, 171; ghiesbreghtiana 21
 Calyptronoma 171; rivalis 170, 171
 Carpentaria acuminata 84, 104
 Caryota 7, 8, 9, 10, 13, 15, 16, 17, 49, 95, 131, 182;
 cumingii 20, 23; mitis 6; urens 181
 Catoblastus 55; praemorsus 73, 74, 75
 Ceroxylon 55, 57, 59, 60, 62, 63, 98; alpinum 55; flex-
 uosum 60; quinduense 55, 60, 62; interruptum 79, 80;
 kloppstockia 73, 79; mooreanum 60, 62; utile 55, 56,
 57, 58
 Chamaedorea costaricana 127; elegans 6, 8, 9, 12; tepejilo-
 te 21
 Chamaerops 144
 Chelyocarpus 120
 Chrysalidocarpus lutescens 6, 10, 31, 33, 34, 35, 36, 37,
 38, 39, 45, 59, 119, 181; madagascariensis 127
 Classified 19, 72, 126
 Coccothrinax 169, 171; alta 170; barbadensis 79, 80; cri-
 nata 19, 72, 167
 Cocos 47; nucifera 20, 21, 22, 23, 49, 59, 90, 172, 180,
 181, 182, 184
 Copernicia cerifera 103; prunifera 103
 Corypha 21, 22, 23, 131; elata 20, 21, 23
 Cryosophila 21, 103, 120, 125; albida 120, 121, 122, 124,
 125; guagara 120; nana 21; warscewiczii 120
 Cyrtostachys lakka 103; renda 49, 103
 Darian, M. E. 152
 Dengler, Nancy G. and Ronald E. Dengler
 Formation of plications in the pinnate leaves of *Chrysalidocarpus lutescens* and the palmate leaves of *Rhapis excelsa* 31
 Dictyocaryum 55, 63; platysepalum 61, 62
 Dictyosperma 131
 Dolby, Warren, J.
 Palms in the microclimates of the San Francisco Bay
 area 92
 Donselman, Henry M., as coauthor 90
 Dransfield, John, as coauthor 3
 Review of *Bombery* and *Rodd*, Palms 102
 Dransfield, John and Natalie W. Uhl
 A magnificent new palm from Madagascar 151
 Halmooorea, a new genus from Madagascar with notes on
Sindroa and *Orania* 163
 Elaeis 47, 127; guineensis 155, 180, 181, 182, 184, 185
 Eugeissona tristis 88; utilis 131, 146
 Euterpe 55, 57, 58, 63, 127, 131; acuminata 73, 80, 81;
 globosa 171; macrospadix 129, 130; precatorea 81
 Euterpe aff. microcarpa 60, 61, 62
 Financial statement 191
 Fullington, J. Garrin, as coauthor 55
 Guassia 171; attenuata 170; princeps 171
 Genera Palmarum 100, 147
 Geonoma 55, 56, 58, 75, 171; euterpoidea 62; pulcherrima
 62; simplicifrons 81; spinescens 81, 82; undata 82, 83
 Ghose, Monoranjan
 Vessel elements in roots of young palms 179
 Gronophyllum ramsayii 19, 72, 167
 Guzman-Rivas, Pablo
 Coconut and other palm use in Mexico and the Philippines
 20
 Halmooorea 163-167; trispatha 166
 Hedyscepe canterburyana 49
 Henderson, Andrew
 Observations on pollination of *Cryosophila albica* 120
 The native palms of Puerto Rico 168
 Heterospatha 103; elata 20, 84, 104
 Hobbs, Horace O. Jr.
 Palm adventures in Costa Rica 128
 House, Alan P. N.
 The ecology of *Oncosperma horridum* on Siberut Island,
 Indonesia 85
 Howea 7, 8, 9, 10, 11, 13, 49, 92, 95; forsteriana 6, 95
 Hydrastele 100, 125; wendlandiana 99
 Hyophorbe 131; indica 130
 Hyospatha pittieri 75, 76, 83
 Hyphaene 103; compressa 103; coriacea 103; dichotoma
 180, 181; natalensis 103
 Iguanura wallichiana 176

- Iriarteae 130, 131; gigantea 130
 Johannesteijsmannia 173, 178; altifrons 173, 178; magnifica 177
 Johnson, Dennis V., as a translator 20
 Notes on the palms of Guinea-Bissau 155
 A bibliography of graduate theses on palms, Part II 186
 Jubaea 95, 96; chilensis 19, 55, 96
 Jubaeopsis caffra 49, 101
 Laccospadix australasica 99, 100
 Laccosperma secundiflorum 161
 Latania loddigesii 19, 72, 167
 Letters 119, 172
 Licuala 3, 131, 172; exigua 171; flabellum 103; gracilis 103; longipes 177; ramsayi 100; spinosa 176
 Linospadix 100; microcarya 100; minor 99, 100
 Livistona 3, 16, 144, 172, 182, 183, 185; australis 99; exigua 1, 3, 4, 5; rotundifolia 20, 180, 181; saribus 21, 127; speciosa 176
 Lodoicea maldivica 49
 Manicaria saccifera 64
 Marojejya darianii 148, 149-154; insignis 151, 153, 154
 Metroxylon 176; sagu 21, 23
 Moratia cerifera 131, 146
 Morgan, P. R. and John Dransfield
 Livistona exigua, a rare Bornean palm refound 3
 Motiska, Dale, Garrin J. Fullington, and Cynthia Anderson
 The renewed quest for high elevation palms 55
 Myers, Ronald L.
 Growth form, growth characteristics and phenology of *Raphia taedigera* in Costa Rican palm swamps 64
 Nannorrhops 95
 Nenga 131
 Neodypsis 49, 103, 131; decaryi 19, 72, 167
 Neonicholsonia 131
 News 49, 99, 144; from Arizona 144; from Southern California 49, 144; from Texas 144
 Normanbya 49, 131
 Nypa 21, 22, 23; fruticans 21, 23, 132, 133
 Oenocarpus 131
 Oncosperma 85; fasciculatum 85; gracilipes 85; horridum 85, 86, 87, 88, 89; platyphyllum 85; tigillarum 85, 87
 Orania 163; appendiculata 99, 163; longisquama 164; sylvicola 176
 Orbignya 21; cohune 21, 22
 Paivoke, Aira E. A.
 Tapping patterns in the Nipa palm (*Nypa fruticans* Wurm) 132
 Palm conferences 101
 Palm literature 102, 162, 186
 Parajubaea coccoides 55
 Phoenix canariensis 55, 56, 95, 96, 97, 144; dactylifera 21; laouireii 21; pusilla 181, 182, 183, 184, 185; reclinata 144, 180, 181, 182, 184, 185; roebelenii 6, 11, 12, 14, 95; rupicola 19, 72, 97, 167, 180, 181, 182; sylvestris 181, 182, 183
 Pholidostachys 171
 Pigafetta 131, filaris 131, 146
 Pinanga 49, 131, 176; sp. 176, 177
 Prestoea montana 171
 Prestoea 131; longipetiolata 83
 Pritchardia 49
 Pseudophoenix sargentii 171
 Ptychosperma 49, 131; elegans 100
 Raphia 66, 67, 68, 69, 70, 71, 72, 131; palma-pinus 161; sudanica 161; taedigera 64, 65, 66, 67
 Raulerson, Charles, obituary 49
 Reinhardtia 131
 Revolving Publications Fund 52
 Rhipidophyllum 125; hystrix 120, 144
 Rhapis 19, 40, 45, 47, 72, 126, 131; excelsa 31, 33, 40, 41, 42, 43, 44, 45, 46, 167
 Rhopaloblaste 103
 Rhopalostylis 49; baueri 103
 Roystonea 49, 59, 127, 168, 170, 182; borinquena 168; regia 19, 90, 181, 185
 Ruhrland, E., letter 172
 Sabal 21, 29, 95, 125, 131; causarium 169; dugesii 19; mexicana 21; palmetto 90, 120, 144
 Salacca edulis 103; zalacca 103, 180, 181, 184
 Scheelea 23, 74, 128; butyracea 23, 82, 83; macrolepis 83, 84; rostrata 128
 Schippia concolor 131, 146
 Sindroa 163; longisquama 163, 164
 Smith, Dent 50
 Sneed, Melvin W.
 Palms in Nepal near the top of the World 15
 Where's Joey? 173
 Socratea 131; durissima 131
 Southern California Chapter 49
 Syagrus sancona 59, 84
 Synechanthus 131; warszewiczianus 127
 Teysmannia 173
 Thrinax 125, 171; morrisii 169, 170; parviflora 120
 Trachycarpus 15, 92, 131; fortunei 55, 127, 144
 Trithrinax 95
 Uhl, Natalie W., as coauthor 151, 613
 Veitchia 179, 185; merrillii 21, 181, 182, 185
 Velez, Ralph, letter from 119
 Verschaffeltia splendida 175
 Walska, Mme. Ganna, obituary notice 144
 Washingtonia 92, 94, 95, 96, 97; filifera 21, 144; robusta 21, 55, 144
 Welfia 171
 Wettinia 55; fascicularis 59, 62
 Wilcox, Merrill 50
 Zambia 103

Principes, 28(4), 1984, p. 191

THE INTERNATIONAL PALM SOCIETY, INC.
STATEMENT OF CASH RECEIPTS
AND DISBURSEMENTS
December 31, 1983

Income

Membership	\$39,765.13	
Interest	2,568.89	
Subscribers	1,569.30	
Seed Bank	3,144.00	
Publications	7,327.10	
Postage	851.50	
Miscellaneous	<u>2,200.55</u>	
Total Income		\$57,426.47

Disbursements

Printing of Catalog	\$32,784.03	
Postage	270.00	
Accounting	150.00	
Travel	995.20	
Bank Charges	27.89	
Rent & Supplies	243.15	
Dues & Subscriptions	129.00	
Miscellaneous	1,450.86	
Grants	<u>6,653.00</u>	
Total Expenses		<u>42,703.13</u>
Excess Receipts over Disbursements		<u><u>\$14,723.34</u></u>


BALANCE SHEET
December 31, 1983

Assets

Petty Cash	\$ 500.00	
"Super Now"	1,524.60	
Coral Gables Federal Savings & Loan	10,061.41	
"IMF"	8,717.36	
First National Bank of Miami	814.59	
Douglas County Bank	15,631.33	
World Savings & Loan	<u>14,947.30</u>	
Total Assets		<u><u>\$52,196.59</u></u>

Fund Balance

Fund Balance—December 31, 1983	\$37,473.25	
Net Receipts over Disbursements	<u>14,723.34</u>	
Balance		<u><u>\$52,196.59</u></u>



GENERAL PALMARUM FUND

SOARS!

DOUBLE YOUR GIVING POWER

Do you work for a company with a Matching Gift Program? Many employers will match your gifts to non-profit organizations dollar for dollar, thus doubling your

gift. If your company does this, your \$100 gift to the International Palm Society would grow to \$200! In most cases, all you have to do is get a Matching Gift Form from your personnel office and fill it out according to your company's instructions. This would mean a great deal to the Society, helping us reach our goals of promoting scientific knowledge on palms.

A list of companies with matching Gift Programs follows, if your employer is not listed, please check with your personnel office.

AT&T	Honeywell
Atlantic Richfield	IBM
Bank America	Montgomery Ward
Coca Cola	Philip Morris
Del Monte	Textron
B. F. Goodrich	U. S. Air

SEND DONATIONS TO:

The International Palm Society, Inc.
for the Revolving Publications Fund,
3616 Mound Ave., Bentura, CA 93003 USA

All donations to The Int. Palm Society, Inc.
are tax deductible in the USA